

Minimal Scenarios for Leptogenesis and CP Violation

G. C. Branco,^{1,*} R. González Felipe,^{1,†} F. R. Joaquim,^{1,‡}

I. Masina,^{2,§} M. N. Rebelo,^{1,¶} and C. A. Savoy^{2,**}

¹*Departamento de Física and Grupo Teórico de Física de Partículas,
Instituto Superior Técnico,
Av. Rovisco Pais, 1049-001 Lisboa, Portugal*

²*Service de Physique Théorique, Laboratoire de la Direction des
Sciences de la Matière du Commissariat à l'Énergie Atomique et
Unité de Recherche Associée au CNRS (URA 2306), CEA-Saclay,
F-91191 Gif-sur-Yvette, France*

Abstract

The relation between leptogenesis and CP violation at low energies is analyzed in detail in the framework of the minimal seesaw mechanism. Working, without loss of generality, in a weak basis where both the charged lepton and the right-handed Majorana mass matrices are diagonal and real, we consider a convenient generic parametrization of the Dirac neutrino Yukawa coupling matrix and identify the necessary condition which has to be satisfied in order to establish a direct link between leptogenesis and CP violation at low energies. In the context of the LMA solution of the solar neutrino problem, we present minimal scenarios which allow for the full determination of the cosmological baryon asymmetry and the strength of CP violation in neutrino oscillations. Some specific realizations of these minimal scenarios are considered. The question of the relative sign between the baryon asymmetry and CP violation at low energies is also discussed.

PACS numbers: 13.35.Hb, 14.60.St, 14.60.Pq, 11.30.Er

Keywords: Leptogenesis, Neutrino masses and mixing, CP violation.

*Electronic address: gbranco@alfa.ist.utl.pt

†Electronic address: gonzalez@gtae3.ist.utl.pt

‡Electronic address: filipe@gtae3.ist.utl.pt

§Electronic address: masina@spht.saclay.cea.fr

¶Electronic address: rebelo@alfa.ist.utl.pt

**Electronic address: savoy@spht.saclay.cea.fr

I. INTRODUCTION

One of the most exciting recent developments in particle physics is the discovery of neutrino oscillations pointed out by the Super-Kamiokande experiment [1] and confirmed by the Sudbury Neutrino Observatory [2]. Neutrino oscillations provide evidence for non-vanishing neutrino masses and mixings, with the novel feature that large leptonic mixing angles are required, in contrast to what happens in the quark sector. Indeed, the combined results from these experiments suggest that, in addition to the large mixing angle required by the atmospheric neutrino data, another large angle should be present in the leptonic sector. This leads to the so-called large mixing angle (LMA) solution of the solar neutrino problem which turns out to be presently the most favored scenario for the explanation of the solar neutrino deficit. From a theoretical point of view, understanding the large leptonic mixing is still an unresolved mystery for which a considerable number of solutions have been proposed [3]. On the other hand, the appearance of neutrino masses much smaller than those of charged leptons is elegantly explained through the seesaw mechanism [4] which can be implemented by extending the standard model (SM) particle content with right-handed neutrinos. These can be easily accommodated in grand unified theories (GUT) where they appear on equal footing with the other SM particles.

The heavy singlet neutrino states can also play an important role in cosmology, namely, in the explanation of the observed cosmological baryon asymmetry. During the last few years, the data collected from the acoustic peaks in the cosmic microwave background radiation [5] has allowed to obtain a precise measurement of the baryon asymmetry of the universe (BAU). The MAP experiment [6] and the PLANCK satellite [7] planned for the near future should further improve this result. At the present time, the measurement of the baryon-to-entropy ratio $Y_B = n_B/s$ is

$$0.7 \times 10^{-10} \lesssim Y_B \lesssim 1.0 \times 10^{-10}. \quad (1)$$

Leptogenesis is one of the most attractive mechanisms to generate the BAU. As first suggested by Fukugita and Yanagida [8], the key ingredient in leptogenesis are the heavy Majorana neutrinos which, once included in the SM, can give rise to a primordial lepton asymmetry through their out-of-equilibrium decays. This lepton asymmetry is subsequently reprocessed into a net baryon asymmetry by the anomalous sphaleron processes.

In spite of being attractive and successful, leptogenesis turns out to be extremely difficult or even impossible to test experimentally in a direct way. This difficulty is obviously related to the large masses of the heavy Majorana neutrino singlets. Nevertheless, the joint analysis of leptogenesis and low-energy neutrino phenomenology can be viewed as an indirect way of testing it and here the experimental results from neutrino oscillation experiments such as those related to the search of leptonic CP violation in the future long-baseline neutrino experiments are extremely valuable [9].

In this paper, we will address the question of linking the amount and sign of the BAU to low-energy neutrino experiments, namely to the sign and strength of the CP asymmetries measured through neutrino oscillations. Our analysis is performed in the weak basis (WB) where the charged lepton mass matrix m_ℓ and the right-handed Majorana matrix M_R are both real and diagonal. In this WB, all CP-violating phases are contained in the Dirac neutrino mass matrix m_D . The matrix m_D is arbitrary and complex, but since three of its nine phases can be eliminated through rephasing, one is left with six independent physical CP-violating phases. In order to study the link between the BAU generated through leptogenesis and CP violation at low energies, it is crucial to use a convenient parametrization of m_D . We shall make use of the fact that any arbitrary complex matrix can, without loss of generality, be written as the product of a unitary matrix U and a lower triangular matrix Y_Δ . We show that U contains three phases which do not contribute to leptogenesis, while the other three phases contained in Y_Δ contribute to both leptogenesis and low-energy CP violation. As a result, a necessary condition for having a link between leptogenesis and low-energy CP breaking is that the matrix U contains no phases, the simplest choice being obviously $U = \mathbb{1}$. Within this class of Dirac neutrino mass matrices, we perform a search of the minimal scenarios where not only a good fit of low-energy neutrino data is obtained but also a link between the observed size and sign of the BAU and the strength of CP violation observable at low energies through neutrino oscillations can be established.

II. GENERAL FRAMEWORK

We work in the framework of a minimal extension of the SM which consists of adding to the standard spectrum one right-handed neutrino per generation. Before gauge symmetry

breaking, the leptonic couplings to the SM Higgs doublet ϕ can be written as:

$$\mathcal{L}_Y = -Y_\nu (\bar{\ell}_L^0, \bar{\nu}_L^0) \tilde{\phi} \nu_R^0 - Y_\ell (\bar{\ell}_L^0, \bar{\nu}_L^0) \phi \ell_R^0 + \text{H.c.}, \quad (2)$$

where $\tilde{\phi} = i\tau_2 \phi^*$. After spontaneous gauge symmetry breaking the leptonic mass terms are given by

$$\begin{aligned} \mathcal{L}_m &= - \left[\bar{\nu}_L^0 m_D \nu_R^0 + \frac{1}{2} \nu_R^{0T} C M_R \nu_R^0 + \bar{\ell}_L^0 m_\ell \ell_R^0 \right] + \text{H.c.} \\ &= - \left[\frac{1}{2} n_L^T C \mathcal{M}^* n_L + \bar{\ell}_L^0 m_\ell \ell_R^0 \right] + \text{H.c.}, \end{aligned} \quad (3)$$

where $m_D = v Y_\nu$ is the Dirac neutrino mass matrix with $v = \langle \phi^0 \rangle / \sqrt{2} \simeq 174 \text{ GeV}$, M_R and $m_\ell = v Y_\ell$ denote the right-handed Majorana neutrino and charged lepton mass matrices, respectively, and $n_L = (\nu_L^0, (\nu_R^0)^c)$. Among all the terms, only the right-handed neutrino Majorana mass term is $\text{SU}(2) \times \text{U}(1)$ invariant and, as a result, the typical scale of M_R can be much above the electroweak symmetry breaking scale v , thus leading to naturally small left-handed Majorana neutrino masses of the order m_D^2/M_R through the seesaw mechanism. In terms of weak-basis eigenstates the leptonic charged current interactions are given by:

$$\mathcal{L}_W = -\frac{g}{\sqrt{2}} W_\mu^+ \bar{\nu}_L^0 \gamma^\mu \ell_L^0 + \text{H.c.}. \quad (4)$$

It is clear from Eqs. (3) and (4) that it is possible to choose, without loss of generality, a weak basis (WB) where both m_ℓ and M_R are diagonal, real and positive. Note that in this WB, m_D is a general complex matrix which contains all the information on CP-violating phases. Since in the present framework there is no $\Delta L = 2$ mass term of the form $\frac{1}{2} \nu_L^{0T} C M_L \nu_L^0$, the total number of CP-violating phases for n generations is given by $n(n-1)$ [10] which are all contained in m_D in this special weak basis¹.

We recall that the full 6×6 neutrino mass matrix $\mathcal{M}$ is diagonalized via the transformation:

$$V^T \mathcal{M}^* V = \mathcal{D}, \quad (5)$$

where $\mathcal{D} = \text{diag}(m_1, m_2, m_3, M_1, M_2, M_3)$, with m_i and M_i denoting the physical masses of the light and heavy Majorana neutrinos, respectively. It is convenient to write V and $\mathcal{D}$ in

¹ The counting of independent CP-violating phases for the general case, where besides m_D and M_R there is also a left-handed Majorana mass term at tree level has been discussed in Ref. [11].

the following form, together with the definition of $\mathcal{M}$:

$$V = \begin{pmatrix} K & Q \\ S & T \end{pmatrix} , \quad \mathcal{D} = \begin{pmatrix} d_\nu & 0 \\ 0 & D_R \end{pmatrix} , \quad \mathcal{M} = \begin{pmatrix} 0 & m_D \\ m_D^T & M_R \end{pmatrix} . \quad (6)$$

From Eq. (5) one obtains, to an excellent approximation, the seesaw formula:

$$d_\nu \simeq -K^\dagger m_D M_R^{-1} m_D^T K^* \equiv K^\dagger \mathcal{M}_\nu K^* , \quad (7)$$

where $\mathcal{M}_\nu$ is the usual light neutrino effective mass matrix. The leptonic charged-current interactions are given by

$$-\frac{g}{\sqrt{2}} (\bar{\ell}_L \gamma_\mu K \nu_L + \bar{\ell}_L \gamma_\mu Q N_L) W^\mu + \text{H.c.} , \quad (8)$$

where ν_i and N_i denote the light and heavy neutrino mass eigenstates, respectively. The matrix K which contains all information on mixing and CP violation at low energies can then be parametrized, after eliminating the unphysical phases, by $K = U_\delta P$ with $P = \text{diag}(1, e^{i\alpha}, e^{i\beta})$ (α and β are Majorana phases) and U_δ a unitary matrix which contains only one (Dirac-type) phase δ . In the limit where the heavy neutrinos exactly decouple from the theory, the matrix K is usually referred as the Pontecorvo-Maki-Nakagawa-Sakata mixing matrix, which from now on we shall denote as U_ν .

CP violation in neutrino oscillations

It has been shown [12] that the strength of CP violation at low energies, observable for example through neutrino oscillations, can be obtained from the following low-energy WB invariant:

$$\mathcal{T}_{CP} = \text{Tr} [\mathcal{H}_\nu, H_\ell]^3 = 6 i \Delta_{21} \Delta_{32} \Delta_{31} \text{Im} [(\mathcal{H}_\nu)_{12}(\mathcal{H}_\nu)_{23}(\mathcal{H}_\nu)_{31}] , \quad (9)$$

where $\mathcal{H}_\nu = \mathcal{M}_\nu \mathcal{M}_\nu^\dagger$, $H_\ell = m_\ell m_\ell^\dagger$ and $\Delta_{21} = (m_\mu^2 - m_e^2)$ with analogous expressions for Δ_{31} , Δ_{32} . This relation can be computed in any weak basis. The low-energy invariant (9) is sensitive to the Dirac-type phase δ and vanishes for $\delta = 0$. On the other hand, it does not depend on the Majorana phases α and β appearing in the leptonic mixing matrix. The quantity $\mathcal{T}_{CP}$ can be fully written in terms of physical observables once

$$\text{Im} [(\mathcal{H}_\nu)_{12}(\mathcal{H}_\nu)_{23}(\mathcal{H}_\nu)_{31}] = -\Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2 \mathcal{J}_{CP} , \quad (10)$$

where the Δm_{ij}^2 's are the usual light neutrino mass squared differences and $\mathcal{J}_{CP}$ is the imaginary part of an invariant quartet appearing in the difference of the CP-conjugated neutrino oscillation probabilities $P(\nu_e \rightarrow \nu_\mu) - P(\bar{\nu}_e \rightarrow \bar{\nu}_\mu)$. One can easily get:

$$\begin{aligned}\mathcal{J}_{CP} &\equiv \text{Im} [(U_\nu)_{11}(U_\nu)_{22}(U_\nu)_{12}^*(U_\nu)_{21}^*] \\ &= \frac{1}{8} \sin(2\theta_{12}) \sin(2\theta_{13}) \sin(2\theta_{23}) \cos(\theta_{13}) \sin \delta ,\end{aligned}\tag{11}$$

where the θ_{ij} are the mixing angles appearing in the standard parametrization adopted in [13]. Alternatively, one can use Eq. (10) and write:

$$\mathcal{J}_{CP} = -\frac{\text{Im} [(\mathcal{H}_\nu)_{12}(\mathcal{H}_\nu)_{23}(\mathcal{H}_\nu)_{31}]}{\Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2} .\tag{12}$$

This expression has the advantage of allowing the computation of the low-energy CP invariant without resorting to the mixing matrix U_ν .

It is also possible to write WB invariants useful to leptogenesis [12] as well as WB invariant conditions for CP conservation in the leptonic sector relevant in specific frameworks [11, 14].

III. CP ASYMMETRIES IN HEAVY MAJORANA NEUTRINO DECAYS

The starting point in leptogenesis scenarios is the CP asymmetry generated through the interference between tree-level and one-loop heavy Majorana neutrino decay diagrams. In the simplest extension of the SM, such diagrams correspond to the decay of the Majorana neutrino into a lepton and a Higgs boson. Considering the decay of one heavy Majorana neutrino N_j , this asymmetry is given by:

$$\varepsilon_j = \frac{\Gamma(N_j \rightarrow \ell \phi) - \Gamma(N_j \rightarrow \bar{\ell} \phi^\dagger)}{\Gamma(N_j \rightarrow \ell \phi) + \Gamma(N_j \rightarrow \bar{\ell} \phi^\dagger)} .\tag{13}$$

In terms of the Dirac neutrino Yukawa couplings the CP asymmetry (13) is [15]:

$$\varepsilon_j = \frac{1}{8\pi(Y_\nu^\dagger Y_\nu)_{jj}} \sum_{k \neq j} \text{Im}[(Y_\nu^\dagger Y_\nu)_{jk}^2] f\left(\frac{M_k^2}{M_j^2}\right) ,\tag{14}$$

where the index j is not summed over in $(Y_\nu^\dagger Y_\nu)_{jj}$. The loop function $f(x)$ includes the one-loop vertex and self-energy corrections to the heavy neutrino decay amplitudes,

$$f(x) = \sqrt{x} \left[(1+x) \ln \left(\frac{x}{1+x} \right) + \frac{2-x}{1-x} \right] .\tag{15}$$

From Eq. (14) it can be readily seen that the CP asymmetries are only sensitive to the CP-violating phases appearing in $Y_\nu^\dagger Y_\nu$ (or equivalently in $m_D^\dagger m_D$) in the WB where M_R and m_ℓ are diagonal.

A. Hierarchical case: $M_1 < M_2 \ll M_3$

In the hierarchical case $M_1 < M_2 \ll M_3$, only the decay of the lightest heavy neutrino N_1 is relevant for leptogenesis, provided the interactions of N_1 are in thermal equilibrium at the time $N_{2,3}$ decay, so that the asymmetries produced by the latter are erased before N_1 decays. In this situation, it is sufficient to take into account the CP asymmetry ε_1 . Since in the limit $x \gg 1$ the function $f(x)$ can be approximated by² $f(x) \simeq -3/(2\sqrt{x})$, we have from Eq. (14)

$$\varepsilon_1 = -\frac{3}{16\pi(Y_\nu^\dagger Y_\nu)_{11}} \sum_{k=2,3} \text{Im}[(Y_\nu^\dagger Y_\nu)_{1k}^2] \frac{M_1}{M_k}, \quad (16)$$

which can be recast in the form [16]

$$\varepsilon_1 \simeq -\frac{3 M_1}{16 \pi} \frac{\text{Im} [Y_\nu^\dagger Y_\nu D_R^{-1} Y_\nu^T Y_\nu^*]_{11}}{(Y_\nu^\dagger Y_\nu)_{11}} = \frac{3 M_1}{16 \pi v^2} \frac{\text{Im} [Y_\nu^\dagger \mathcal{M}_\nu Y_\nu^*]_{11}}{(Y_\nu^\dagger Y_\nu)_{11}}, \quad (17)$$

using the seesaw relation given in Eq. (7).

B. Two-fold quasi-degeneracy: $M_1 \simeq M_2 \ll M_3$

In the context of thermal leptogenesis, when the observed baryon asymmetry is generated through the decays of the lightest heavy Majorana neutrino N_1 , there exists an upper bound on the CP asymmetry ε_1 which directly depends on the mass of the lightest neutrino M_1 [17]. In turn, such a bound implies a lower bound on the lightest mass M_1 , typically $M_1 \gtrsim 10^8$ GeV. The latter bound³ is however barely compatible with the reheating temperature bound $T_R \lesssim 10^8 \sim 10^9$ GeV required in several supergravity models in order to avoid a gravitino overproduction [19]. To overcome this problem⁴ one can consider, for instance, the decays of two heavy neutrinos which are quasi-degenerate in mass, $M_1 \simeq M_2$. In this case, the CP asymmetries ε_i are enhanced due to self-energy contributions [21] and the required baryon asymmetry can be produced by right-handed heavy neutrinos with moderate masses

² This approximation can be reasonably used for $x \gtrsim 15$.

³ A more stringent constraint, $M_1 \gtrsim 10^{10}$ GeV, is obtained in [18].

⁴ Another possible solution to the gravitino problem is to consider non-thermal production mechanisms [20]. Since in these cases the condition $M_1 < T_R$ is not required, the gravitino problem is easily avoided once heavy particles can be created with a relatively low reheating temperature without threatening big bang nucleosynthesis.

$M_1 \simeq M_2 \lesssim 10^8$ GeV. Moreover, it has been shown that in the presence of small Dirac-type leptonic mixing at high energies and GUT-inspired Dirac neutrino Yukawa couplings, the heavy Majorana neutrino degeneracy is compatible with the LMA solar solution [22].

Let us assume that the heavy Majorana neutrinos N_1 and N_2 are quasi-degenerate. It is useful to define the parameter δ_N which represents the degree of degeneracy between the masses M_1 and M_2 as

$$\delta_N = \frac{M_2}{M_1} - 1. \quad (18)$$

Since $M_1 \simeq M_2$, we expect $\delta_N \ll 1$. For the perturbative approach to remain valid, the tree-level decay width Γ_i for each of the heavy Majorana neutrinos must be much smaller than the mass difference between them. This is translated into the relations

$$\Gamma_i = \frac{(H_\nu)_{ii} M_i}{8\pi} \ll M_2 - M_1 = \delta_N M_1, \quad i = 1, 2, \quad (19)$$

where $H_\nu = Y_\nu^\dagger Y_\nu$. From this equation we can find the following lower bound for δ_N :

$$\delta_N \gg \max \left\{ \frac{(H_\nu)_{ii} M_i}{8\pi M_1} \right\}_{i=1,2}. \quad (20)$$

Assuming that this criterium is verified, the CP asymmetries ε_i can be obtained combining Eqs. (14) and (18). We find

$$\begin{aligned} \varepsilon_1 &= -\frac{1}{8\pi (H_\nu)_{11}} \left\{ \text{Im}[(H_\nu)_{21}^2] f[(1 + \delta_N)^2] - \frac{3}{2} \text{Im}[(H_\nu)_{31}^2] \frac{M_1}{M_3} \right\}, \\ \varepsilon_2 &= -\frac{1}{8\pi (H_\nu)_{22}} \left\{ \text{Im}[(H_\nu)_{12}^2] f[(1 + \delta_N)^{-2}] - \frac{3}{2} \text{Im}[(H_\nu)_{32}^2] \frac{M_2}{M_3} \right\}. \end{aligned} \quad (21)$$

Taking into account that for $\delta_N \ll 1$ the function $f[(1 + \delta_N)^{\pm 2}]$ can be approximated by⁵:

$$f[(1 + \delta_N)^{\pm 2}] \simeq \mp \frac{1}{2\delta_N}, \quad (22)$$

we obtain:

$$\varepsilon_j = \frac{1}{16\pi (H_\nu)_{jj}} \left[\frac{\text{Im}[(H_\nu)_{21}^2]}{\delta_N} + 3 \text{Im}[(H_\nu)_{3j}^2] \frac{M_j}{M_3} \right], \quad j = 1, 2. \quad (23)$$

Typically, the term proportional to M_j/M_3 can be neglected and in this case ε_1 and ε_2 have the same sign. This aspect turns out to be relevant for the discussion on the relative sign between the BAU and low-energy leptonic CP violation.

⁵ For $\delta_N < 10^{-2}$ the error associated to this approximation is less than 1%.

IV. ON THE CONNECTION BETWEEN LEPTOGENESIS AND LOW-ENERGY CP VIOLATION

In this section we analyze the possible link between CP violation at low energies, measurable for example through neutrino oscillations, and leptogenesis. The possibility of such a connection has been previously analyzed in the literature [12, 23, 24]. Nevertheless, we find it worthwhile presenting here a thorough discussion on the subject. In particular, we will address the following questions:

- If the strength of CP violation at low energies in neutrino oscillations is measured, what can one infer about the viability or non-viability of leptogenesis?
- From the sign of the BAU, can one predict the sign of the CP asymmetries at low energies, namely the sign of $\mathcal{J}_{CP}$?

We will show that having an explicit parametrization of m_D (or equivalently of $Y_\nu = m_D/v$) is crucial not only to determine which phases are responsible for leptogenesis and which ones are relevant for leptonic CP violation at low energies, but also to analyze the relationship between these two phenomena.

From the available neutrino oscillation data, one obtains some information on the effective neutrino mass matrix $\mathcal{M}_\nu$ which can be decomposed in the following way:

$$U_\nu d_\nu U_\nu^T = \mathcal{M}_\nu \equiv L L^T, \quad L \equiv i m_D D_R^{-1/2}. \quad (24)$$

The extraction of L from $\mathcal{M}_\nu$ suffers from an intrinsic ambiguity [25] in the sense that, given a particular solution L_0 of Eq. (24), the matrix $L = L_0 R$ will also satisfy this equation, provided that R is an arbitrary orthogonal complex matrix, $R \in O(3, \mathbb{C})$, i.e. $R R^T = \mathbb{1}$. It is useful to take as a reference solution $L_0 \equiv U_\nu d_\nu^{1/2}$, so that:

$$L \equiv U_\nu d_\nu^{1/2} R. \quad (25)$$

Since three of the phases of m_D can be eliminated, the matrix L has 15 independent parameters. The parametrization of L given in Eq. (25) has the interesting feature that all its parameters are conveniently distributed among U_ν , d_ν and R , which contain 6 (3 angles + 3 phases), 3 and 6 (3 angles + 3 phases) independent parameters, respectively. Of the 18 parameters present in the Lagrangian of the fundamental theory described by m_D and D_R ,

only 9 appear at low energy in $\mathcal{M}_\nu$ through the seesaw mechanism. To further disentangle m_D from D_R in L , one needs the 3 remaining inputs, namely the three heavy Majorana masses of D_R . As for the meaning of the information encoded in R , it turns out that the pattern of this matrix has a suggestive interpretation in terms of the different rôles played by the heavy neutrinos in the seesaw mechanism. In fact, R can be viewed as a dominance matrix [26] since it gives the weights of each heavy Majorana neutrino in the determination of the different light neutrino masses m_i [27]. The fact that R_{ij}^2 are weights for m_i is quite obvious due to the orthogonality of R :

$$m_i = \sum_j m_i R_{ij}^2. \quad (26)$$

On the other hand, since $U_\nu^\dagger m_D = -i d_\nu^{1/2} R D_R^{1/2}$, the single contribution $m_i R_{ij}^2$ is also given by:

$$m_i R_{ij}^2 = -\frac{(U_\nu^\dagger m_D)_{ij}^2}{M_j} \equiv \frac{X_{ij}}{M_j}. \quad (27)$$

Therefore, once U_ν is fixed, each weight R_{ij}^2 just depends on the mass M_j of the j -th heavy Majorana neutrino and on its couplings with the left-handed neutrinos $(m_D)_{kj}$. Thus, the contribution of each heavy neutrino to m_i is well defined and expressed by the weight $\text{Re}(R_{ij}^2)$. One may roughly say that the heavy Majorana neutrino with mass M_j dominates⁶ in m_i if

$$\frac{|\text{Re}(X_{ij})|}{M_j} \gg \frac{|\text{Re}(X_{ik})|}{M_k}, \quad k \neq j \quad (28)$$

which implies $|\text{Re}(R_{ij}^2)| \gg |\text{Re}(R_{ik}^2)|$. So that, if one of the heavy Majorana neutrino gives the dominant contribution to one of the masses m_i , this information is encoded in the structure of R . The interpretation in terms of weights is straightforward for the rotational part of R . However, one has to be careful because in the presence of the three boosts (controlled by the three phases) the weights R_{ij}^2 are not necessarily real and positive. Although this situation is more subtle, the above dominance arguments still hold.

Coming back to the connection between leptogenesis and low-energy data, it is important to note that U_ν does not appear in the relevant combination for leptogenesis $Y_\nu^\dagger Y_\nu$, in the same way as R does not appear in $\mathcal{M}_\nu$. Indeed, one has:

$$m_D^\dagger m_D = D_R^{1/2} R^\dagger d_\nu R D_R^{1/2}. \quad (29)$$

⁶ See for instance Refs. [29, 30] for other approaches to the dominance mechanism.

From the above discussion, it follows that it is possible to write m_D in the form $m_D = -i U_\nu d_\nu^{1/2} R D_R^{1/2}$ in such a way that leptogenesis and the low-energy neutrino data (contained in $\mathcal{M}_\nu$) depend on two independent sets of CP-violating phases, respectively those in R and those in U_ν . In particular, one may have viable leptogenesis even in the limit where there are no CP-violating phases (neither Dirac nor Majorana) in U_ν and hence, no CP violation at low energies [28]. Therefore, in general it is not possible to establish a link between low-energy CP violation and leptogenesis. This connection is model dependent: it can be drawn only by specifying a particular *ansatz* for the fundamental parameters of the seesaw, m_D and D_R , as will be done in the following sections.

The relevance of the matrix R for leptogenesis can be rendered even more explicit [27] by rewriting the ε_1 asymmetry by means of Eq. (29) and defining $R_{ij} = |R_{ij}|e^{i\varphi_{ij}/2}$, $\Delta m_\odot^2 \equiv \Delta m_{21}^2$ and $\Delta m_\oplus^2 \equiv \Delta m_{32}^2$. In the case of hierarchical heavy Majorana neutrinos, say $M_1 \ll M_2 \ll M_3$ one obtains

$$\varepsilon_1 \simeq \frac{3}{16\pi} \frac{M_1}{v^2} \frac{\Delta m_\oplus^2 |R_{31}|^2 \sin \varphi_{31} - \Delta m_\odot^2 |R_{11}|^2 \sin \varphi_{11}}{m_1 |R_{11}|^2 + m_2 |R_{21}|^2 + m_3 |R_{31}|^2}, \quad (30)$$

and we recover what one would have expected by intuition, namely that the physical quantities involved in determining ε_1 are just M_1 , the spectrum of the light neutrinos, m_i , and the first column of R , which expresses the composition of the lightest heavy Majorana neutrino in terms of the light neutrino masses m_i . In the case $M_1 \simeq M_2 \ll M_3$, similar expressions hold,

$$\varepsilon_j \simeq \frac{1}{16\pi v^2} \frac{M_1 M_2}{M_2 - M_1} \frac{\text{Im}[(R^\dagger d_\nu R)_{21}^2]}{(R^\dagger d_\nu R)_{jj}}, \quad j = 1, 2. \quad (31)$$

where now also the mass M_2 and the second column of R are involved. A detailed study of the relevance of the matrix R for leptogenesis is under way [31].

As stressed before, different *ansätze* for R have no direct impact on CP violation at low energy; the impact is in a sense indirect because R specifies if dominance of some heavy Majorana neutrino is at work in the seesaw mechanism [26].

In conclusion, the link between leptogenesis and low-energy CP violation can only be established in the framework of specific *ansätze* for the leptonic mass terms of the Lagrangian. We shall derive a necessary condition for such a link to exist. In order to obtain this connection, it is convenient to use a triangular parametrization for m_D , which we describe next.

Triangular parametrization

It can be easily shown that any arbitrary complex matrix can be written as the product of a unitary matrix U with a lower triangular matrix Y_Δ . In particular, the Dirac neutrino mass matrix can be written as:

$$m_D = v U Y_\Delta, \quad (32)$$

with Y_Δ of the form:

$$Y_\Delta = \begin{pmatrix} y_{11} & 0 & 0 \\ y_{21} e^{i\phi_{21}} & y_{22} & 0 \\ y_{31} e^{i\phi_{31}} & y_{32} e^{i\phi_{32}} & y_{33} \end{pmatrix}, \quad (33)$$

where y_{ij} are real positive numbers. Since U is unitary, in general it contains six phases. However, three of these phases can be rephased away by a simultaneous phase transformation on ν_L^0, ℓ_L^0 , which leaves the leptonic charged current invariant. Under this transformation, $m_D \rightarrow P_\xi m_D$, with $P_\xi = \text{diag}(e^{i\xi_1}, e^{i\xi_2}, e^{i\xi_3})$. Furthermore, Y_Δ defined in Eq. (33) can be written as:

$$Y_\Delta = P_\beta^\dagger \hat{Y}_\Delta P_\beta, \quad (34)$$

where $P_\beta = \text{diag}(1, e^{i\beta_1}, e^{i\beta_2})$ with $\beta_1 = -\phi_{21}$, $\beta_2 = -\phi_{31}$ and

$$\hat{Y}_\Delta = \begin{pmatrix} y_{11} & 0 & 0 \\ y_{21} & y_{22} & 0 \\ y_{31} & y_{32} e^{i\sigma} & y_{33} \end{pmatrix}, \quad (35)$$

with $\sigma = \phi_{32} - \phi_{31} + \phi_{21}$. It follows then from Eqs. (32) and (34) that the matrix m_D can be decomposed in the form

$$m_D = v U_\rho P_\alpha \hat{Y}_\Delta P_\beta, \quad (36)$$

where $P_\alpha = \text{diag}(1, e^{i\alpha_1}, e^{i\alpha_2})$ and U_ρ is a unitary matrix containing only one phase ρ . Therefore, in the WB where m_ℓ and M_R are diagonal and real, the phases $\rho, \alpha_1, \alpha_2, \sigma, \beta_1$ and β_2 are the only physical phases characterizing CP violation in the leptonic sector. The phases relevant for leptogenesis are those contained in $m_D^\dagger m_D$. From Eqs. (34)-(36) we

conclude that these phases are σ , β_1 and β_2 , which are linear combinations of the phases ϕ_{ij} . On the other hand, all the six phases of m_D contribute to the three phases of the effective neutrino mass matrix at low energies [12] which in turn controls CP violation in neutrino oscillations. Since the phases α_1 , α_2 and ρ do not contribute to leptogenesis, it is clear that a necessary condition for a direct link between leptogenesis and low-energy CP violation to exist is the requirement that the matrix U in Eq. (32) contains no CP-violating phases. Note that, although the above condition was derived in a specific WB and using the parametrization of Eq. (32), it can be applied to any model. This is due to the fact that starting from arbitrary leptonic mass matrices, one can always make WB transformations to render m_ℓ and M_R diagonal, while m_D has the form of Eq. (32). A specific class of models which satisfy the above necessary condition in a trivial way are those for which $U = \mathbb{1}$, leading to $m_D = v Y_\Delta$. This condition is necessary but not sufficient to allow for a prediction of the sign of the CP asymmetry in neutrino oscillations, given the observed sign of the BAU together with the low-energy data. Therefore, we will consider next a more restrictive class of matrices m_D of this form and we will show that, in an appropriate limiting case, our structures for m_D lead to the ones assumed by Frampton, Glashow and Yanagida in [24].

V. MINIMAL SCENARIOS

From the analysis carried out in the previous section, it becomes clear that the computation of the cosmological baryon asymmetry Y_B in leptogenesis scenarios strongly depends on the Yukawa structure of the Dirac neutrino mass term and on the heavy Majorana neutrino mass spectrum. Moreover, if one assumes that the seesaw mechanism is responsible for the smallness of the neutrino masses, then the connection between the baryon asymmetry and low-energy neutrino physics is unavoidable. In fact, this constitutes an advantage for the leptogenesis mechanism when compared to other baryogenesis scenarios. In the context of supersymmetric extensions of the SM it is possible (although not always simple) to combine the study of leptogenesis and neutrino physics with other physical phenomena like flavor-violating decays [18, 32]. In general, this analysis does not give us definite answers, yet it may help to discriminate among certain neutrino mass models.

Recently, a considerable amount of work has been done aiming at relating viable lepto-

genesis to all the available low-energy neutrino data coming from solar, atmospheric and reactor experiments [33]. Roughly speaking, two different approaches to the problem are to be found. The first one is based on the computation of the baryon asymmetry as a function of the lightest heavy Majorana neutrino mass M_1 , the CP asymmetry ε_1 and the so-called effective neutrino mass $\tilde{m}_1 = (m_D^\dagger m_D)_{11}/M_1$ [34]. By solving the Boltzmann equations, this kind of analysis provides valuable information on the ranges of these parameters that lead to an acceptable value of Y_B . The weak point of this procedure lies on the fact that the input information depends on quantities which are not sensitive to the full structure of Y_ν and M_R and, therefore, no further conclusions can be drawn about the class of models which can lead to acceptable values of the input parameters referred above. In fact, the values of M_1 , $\tilde{m}_1$ and ε_1 should not be taken as independent parameters. The second approach is based upon initial assumptions on the structure of Y_ν and M_R at high energies which are fixed by recurring to theoretical arguments like for example grand unified theories or flavor symmetries. Although in this framework some generality is lost, it has the advantage that one can compute the generated baryon asymmetry and, simultaneously, perform a low-energy neutrino data analysis. It is precisely the latter approach that we shall follow in the present work.

In this section we present a class of minimal scenarios for leptogenesis based on the triangular decomposition of Y_ν given in Eqs. (32) and (33). Namely, we would like to answer the question of how simple can the structure of the Dirac neutrino Yukawa coupling matrix be in order not only to get an acceptable value of the baryon asymmetry but also to accommodate the neutrino data provided by the atmospheric, solar and reactor neutrino experiments. In particular, we require the non-vanishing of the CP asymmetry generated in the decays of the lightest heavy Majorana neutrino, since the final value of the baryon asymmetry crucially depends on this quantity. Throughout our analysis we shall also consider the predictions on the CP-violating effects at low energies.

In the previous section we have seen that the Dirac neutrino Yukawa coupling matrix Y_ν can be decomposed into the product of a unitary matrix U and a lower-triangular matrix Y_Δ (cf. Eq. 32). It was also shown that the CP asymmetries ε_j generated in the heavy Majorana neutrino decays do not depend on the matrix U . In the special case $U = \mathbb{1}$ it is possible to establish the connection between leptogenesis, low-energy CP violation and neutrino mixing, since the same phases affect these phenomena. We classify this scenario

as a minimal scenario for leptogenesis and CP violation in the sense that the CP-violating sources that do not contribute to leptogenesis are neglected. On the other hand, if $U \neq \mathbb{1}$ this connection is not trivial. Therefore, from now on we will consider the case $U = \mathbb{1}$ which implies the following simple structure for the Dirac neutrino mass matrix:

$$m_D = v Y_\Delta = v \begin{pmatrix} y_{11} & 0 & 0 \\ y_{21} e^{i\phi_{21}} & y_{22} & 0 \\ y_{31} e^{i\phi_{31}} & y_{32} e^{i\phi_{32}} & y_{33} \end{pmatrix}. \quad (37)$$

Then, from Eq. (14) the CP asymmetry generated in the decay of the heavy Majorana neutrino N_j is

$$\varepsilon_j = -\frac{1}{8\pi(H_\Delta)_{jj}} \sum_{i \neq j} \text{Im}[(H_\Delta)_{ij}^2] f_{ij}, \quad (38)$$

where

$$H_\Delta = Y_\Delta^\dagger Y_\Delta \quad , \quad f_{ij} = f\left(\frac{M_i^2}{M_j^2}\right), \quad (39)$$

with $f(x)$ defined in Eq. (15).

From Eqs. (37) and (39) we readily obtain

$$\begin{aligned} \text{Im}[(H_\Delta)_{21}^2] &= y_{21}^2 y_{22}^2 \sin(2\phi_{21}) + 2 y_{21} y_{22} y_{31} y_{32} \sin \theta_1 + y_{31}^2 y_{32}^2 \sin \theta_2, \\ \text{Im}[(H_\Delta)_{31}^2] &= y_{31}^2 y_{33}^2 \sin(2\phi_{31}), \\ \text{Im}[(H_\Delta)_{32}^2] &= y_{32}^2 y_{33}^2 \sin(2\phi_{32}), \end{aligned} \quad (40)$$

with $\theta_1 = \phi_{21} + \phi_{31} - \phi_{32}$ and $\theta_2 = 2(\phi_{31} - \phi_{32})$.

All the information about neutrino masses and mixing is fully contained in the effective neutrino mass matrix $\mathcal{M}_\nu$ which is determined through the seesaw formula given by Eq. (7). In this case

$$\mathcal{M}_\nu = \frac{v^2}{M_1} \begin{pmatrix} y_{11}^2 & y_{11}y_{21}e^{i\phi_{21}} & y_{11}y_{31}e^{i\phi_{31}} \\ y_{11}y_{21}e^{i\phi_{21}} & y_{21}^2 e^{2i\phi_{21}} + y_{22}^2 \frac{M_1}{M_2} & y_{21}y_{31}e^{i(\phi_{31}+\phi_{21})} + y_{22}y_{32} \frac{M_1}{M_2} e^{i\phi_{32}} \\ y_{11}y_{31}e^{i\phi_{31}} & y_{21}y_{31}e^{i(\phi_{31}+\phi_{21})} + y_{22}y_{32} \frac{M_1}{M_2} e^{i\phi_{32}} & y_{31}^2 e^{2i\phi_{31}} + y_{33}^2 \frac{M_1}{M_3} + y_{32}^2 \frac{M_1}{M_2} e^{2i\phi_{32}} \end{pmatrix}. \quad (41)$$

It follows from Eqs. (38)-(41) that, in principle, one can obtain simultaneously viable values for the CP asymmetries ε_j and a phenomenologically acceptable effective neutrino mass

matrix in order to reproduce the solar, atmospheric and reactor neutrino data. This can be achieved by consistently choosing the values of the free parameters y_{ij} , M_i and ϕ_{ij} . Yet, a closer look at Eqs. (38)-(40) shows that there are terms contributing to ε_j which vanish independently from the others. This means that a non-vanishing value of ε_j can be guaranteed even for simpler structures for Y_ν , which can be obtained from Y_Δ assuming additional zero entries in the lower triangle⁷. The results are given in Table I where we present the textures constructed from Y_Δ by neglecting one (textures I-III) and two (textures IV-VI) off-diagonal entries. The form of the effective neutrino mass matrix $\mathcal{M}_\nu$ and the expressions for the CP asymmetries $\varepsilon_{1,2}$ for each case are also given⁸.

Let us first discuss textures IV-VI. For these three textures the effective neutrino mass matrix $\mathcal{M}_\nu$ predicts a complete decoupling of one light neutrino from the other two. This is in disagreement with the available neutrino data which indicates that, in the framework of the LMA solution, only one neutrino mixing angle should be small, namely θ_{13} , instead of two as predicted by textures IV-VI. Furthermore, texture VI predicts a vanishing value for the CP asymmetry in the decay of the lightest heavy Majorana neutrino, implying $Y_B = 0$ in the case of hierarchical heavy Majorana neutrinos. Texture III can also be excluded on the grounds that it cannot account for the large solar angle. To illustrate this, let us write for this texture the matrix L given in Eq. (24) as

$$L = i m_D D_R^{-1/2} \equiv \begin{pmatrix} z_1 & 0 & 0 \\ z_2 & y_2 & 0 \\ z_3 & 0 & x_3 \end{pmatrix}, \quad (42)$$

where z_1 , y_2 and x_3 are real and positive [26]. We also take z_2 and z_3 real to simplify the discussion. Considering first the hierarchical case $\Delta m_{\text{atm}}^2 \simeq m_3^2$, a large atmospheric angle and $m_2 < m_3$ naturally arise if $z_2 \sim z_3 > y_2, x_3$. More precisely, $\tan \theta_{23} \simeq z_2/z_3$. In addition, this implies

$$\tan \theta_{13} \simeq \frac{z_1}{\sqrt{z_2^2 + z_3^2}} \equiv t_{13}, \quad (43)$$

⁷ Notice however that the vanishing of diagonal elements in Y_Δ would imply $\det(m_D) = 0$ and consequently, $\det(\mathcal{M}_\nu) = 0$, leading to the existence of massless light neutrinos.

⁸ In commonly used language, textures I-III and IV-VI belong to the classes of four and five texture zero matrices, respectively. For a complete discussion on seesaw realization of texture-zero mass matrices see e.g. [35] and for its implications at low energies see e.g. [36].

TABLE I: Minimal textures based on the triangular decomposition of m_D and their respective light neutrino mass matrix $\mathcal{M}_\nu$ and CP-asymmetries ε_1 and ε_2 .

Texture	Y_Δ	$\frac{M_1}{v^2} \mathcal{M}_\nu$	$\varepsilon_{1,2}$
I	$\begin{pmatrix} y_{11} & 0 & 0 \\ y_{21} e^{i\phi_{21}} & y_{22} & 0 \\ 0 & y_{32} e^{i\phi_{32}} & y_{33} \end{pmatrix}$	$\begin{pmatrix} y_{11}^2 & y_{11} y_{21} e^{i\phi_{21}} & 0 \\ y_{11} y_{21} e^{i\phi_{21}} & y_{21}^2 e^{2i\phi_{21}} + \frac{M_1}{M_2} y_{22}^2 & y_{22} y_{32} \frac{M_1}{M_2} e^{i\phi_{32}} \\ 0 & y_{22} y_{32} \frac{M_1}{M_2} e^{i\phi_{32}} & y_{33}^2 \frac{M_1}{M_3} + y_{32}^2 \frac{M_1}{M_2} e^{2i\phi_{32}} \end{pmatrix}$	$\varepsilon_1^{\text{I}} = -\frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21})}{8\pi (y_{11}^2 + y_{21}^2)} f_{21}$ $\varepsilon_2^{\text{I}} = \frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21}) f_{12} - y_{32}^2 y_{33}^2 \sin(2\phi_{32}) f_{32}}{8\pi (y_{22}^2 + y_{32}^2)}$
II	$\begin{pmatrix} y_{11} & 0 & 0 \\ 0 & y_{22} & 0 \\ y_{31} e^{i\phi_{31}} & y_{32} e^{i\phi_{32}} & y_{33} \end{pmatrix}$	$\begin{pmatrix} y_{11}^2 & 0 & y_{11} y_{31} e^{i\phi_{31}} \\ 0 & y_{22}^2 \frac{M_1}{M_2} & y_{32} y_{22} \frac{M_1}{M_2} e^{i\phi_{32}} \\ y_{11} y_{31} e^{i\phi_{31}} & y_{32} y_{22} \frac{M_1}{M_2} e^{i\phi_{32}} & y_{31}^2 e^{2i\phi_{31}} + y_{33}^2 \frac{M_1}{M_3} + y_{32}^2 \frac{M_1}{M_2} e^{2i\phi_{32}} \end{pmatrix}$	$\varepsilon_1^{\text{II}} = \frac{y_{31}^2 y_{32}^2 \sin[2(\phi_{32} - \phi_{31})] f_{21} - y_{31}^2 y_{33}^2 \sin(2\phi_{31}) f_{31}}{8\pi (y_{11}^2 + y_{31}^2)}$ $\varepsilon_2^{\text{II}} = \frac{y_{31}^2 y_{32}^2 \sin[2(\phi_{31} - \phi_{32})] f_{12} - y_{32}^2 y_{33}^2 \sin(2\phi_{32}) f_{32}}{8\pi (y_{22}^2 + y_{32}^2)}$
III	$\begin{pmatrix} y_{11} & 0 & 0 \\ y_{21} e^{i\phi_{21}} & y_{22} & 0 \\ y_{31} e^{i\phi_{31}} & 0 & y_{33} \end{pmatrix}$	$\begin{pmatrix} y_{11}^2 & y_{11} y_{21} e^{i\phi_{21}} & y_{11} y_{31} e^{i\phi_{31}} \\ y_{11} y_{21} e^{i\phi_{21}} & y_{21}^2 e^{2i\phi_{21}} + y_{22}^2 \frac{M_1}{M_2} & y_{21} y_{31} e^{i(\phi_{21} + \phi_{31})} \\ y_{11} y_{31} e^{i\phi_{31}} & y_{21} y_{31} e^{i(\phi_{21} + \phi_{31})} & y_{31}^2 e^{2i\phi_{31}} + y_{33}^2 \frac{M_1}{M_3} \end{pmatrix}$	$\varepsilon_1^{\text{III}} = -\frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21}) f_{21} + y_{31}^2 y_{33}^2 \sin(2\phi_{31}) f_{31}}{8\pi (y_{11}^2 + y_{21}^2 + y_{31}^2)}$ $\varepsilon_2^{\text{III}} = \frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21})}{8\pi y_{22}^2} f_{12}$
IV	$\begin{pmatrix} y_{11} & 0 & 0 \\ y_{21} e^{i\phi_{21}} & y_{22} & 0 \\ 0 & 0 & y_{33} \end{pmatrix}$	$\begin{pmatrix} y_{11}^2 & y_{11} y_{21} e^{i\phi_{21}} & 0 \\ y_{11} y_{21} e^{i\phi_{21}} & y_{21}^2 e^{2i\phi_{21}} + y_{22}^2 \frac{M_1}{M_2} & 0 \\ 0 & 0 & y_{33}^2 \frac{M_1}{M_3} \end{pmatrix}$	$\varepsilon_1^{\text{IV}} = -\frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21})}{8\pi (y_{11}^2 + y_{21}^2)} f_{21}$ $\varepsilon_2^{\text{IV}} = \frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21})}{8\pi y_{22}^2} f_{12}$
V	$\begin{pmatrix} y_{11} & 0 & 0 \\ 0 & y_{22} & 0 \\ y_{31} e^{i\phi_{31}} & 0 & y_{33} \end{pmatrix}$	$\begin{pmatrix} y_{11}^2 & 0 & y_{11} y_{31} e^{i\phi_{31}} \\ 0 & y_{22}^2 \frac{M_1}{M_2} & 0 \\ y_{11} y_{31} e^{i\phi_{31}} & 0 & y_{31}^2 e^{2i\phi_{31}} + y_{33}^2 \frac{M_1}{M_3} \end{pmatrix}$	$\varepsilon_1^{\text{V}} = -\frac{y_{31}^2 y_{33}^2 \sin(2\phi_{31})}{8\pi (y_{11}^2 + y_{31}^2)} f_{31}$ $\varepsilon_2^{\text{V}} = 0$
VI	$\begin{pmatrix} y_{11} & 0 & 0 \\ 0 & y_{22} & 0 \\ 0 & y_{32} e^{i\phi_{32}} & y_{33} \end{pmatrix}$	$\begin{pmatrix} y_{11}^2 & 0 & 0 \\ 0 & y_{22}^2 \frac{M_1}{M_2} & y_{22} y_{32} \frac{M_1}{M_2} e^{i\phi_{32}} \\ 0 & y_{22} y_{32} \frac{M_1}{M_2} e^{i\phi_{32}} & y_{33}^2 \frac{M_1}{M_3} + y_{32}^2 \frac{M_1}{M_2} e^{2i\phi_{32}} \end{pmatrix}$	$\varepsilon_1^{\text{VI}} = 0$ $\varepsilon_2^{\text{VI}} = -\frac{y_{32}^2 y_{33}^2 \sin(2\phi_{32})}{8\pi (y_{22}^2 + y_{32}^2)} f_{32}$

which has to satisfy the CHOOZ bound, $t_{13} \leq 0.2$ [37]. The solar angle is approximately given by

$$\tan(2\theta_{12}) \simeq 2 t_{13} \frac{y_2^2 + x_3^2}{\frac{z_3}{z_2} y_2^2 + \frac{z_2}{z_3} x_3^2 - t_{13}(\frac{z_2}{z_3} y_2^2 + \frac{z_3}{z_2} x_3^2)} \leq \frac{2 t_{13} \beta}{1 - t_{13} \beta^2} \equiv B_{12}(\beta, t_{13}), \quad (44)$$

where $\beta \equiv \max(\tan \theta_{23}, \cot \theta_{23})$. The upper bound $B_{12}(\beta, t_{13} = 0.2)$ is an increasing function of β . Experimentally $0.7 \leq \beta \leq 1.7$, so that $B_{12}^{\max} = B_{12}(1.7, 0.2) = 4$. This corresponds to $\tan^2 \theta_{12} \leq 0.3$, while the fitted LMA solution requires $\tan^2 \theta_{12} > 0.3$. We thus conclude that type III cannot account for the observed large solar angle in the hierarchical case. Moreover, it turns out that it cannot also account for the pattern of $\mathcal{M}_\nu$ required by inverse hierarchy since, in this case, the atmospheric oscillation fit would require $z_1 z_2 \simeq z_1 z_3 \simeq \sqrt{\Delta m_{\text{atm}}^2}$ and $z_1^2, z_2^2, z_3^2 \ll \sqrt{\Delta m_{\text{atm}}^2}$. Finally, the degenerate case can be accommodated only by tuning the elements of L . In view of this, from now on we will focus our analysis only on textures I and II.

Another interesting fact which comes out from the observation of Table I is that the phase content for textures I-II can be further reduced. Indeed, it is straightforward to show that with only one phase in Y_Δ it is possible to obtain a non-vanishing CP asymmetry ε_1 .

As it has been stated in Section II, the strength of CP violation at low energies is sensitive to the value of the CP invariant $\mathcal{J}_{CP}$ given by Eq. (12). For the textures I and II we get:

$$\begin{aligned} \mathcal{J}_{CP}^{\text{I}} &= \frac{y_{11}^2 y_{21}^2 y_{32}^2 y_{22}^2 v^{12}}{M_1^3 M_2^3 \Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2} \left[(y_{21}^2 y_{32}^2 + y_{11}^2 y_{22}^2 + y_{11}^2 y_{32}^2) \sin(2\phi_{21}) - y_{22}^2 y_{33}^2 \frac{M_1}{M_3} \sin(2\phi_{32}) \right. \\ &\quad \left. + y_{33}^2 (y_{11}^2 + y_{21}^2) \frac{M_2}{M_3} \sin[2(\phi_{21} - \phi_{32})] \right], \\ \mathcal{J}_{CP}^{\text{II}} &= \frac{y_{11}^2 y_{22}^2 y_{31}^2 y_{32}^2 v^{12}}{M_1^3 M_2^3 \Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2} \left[(y_{22}^2 y_{31}^2 + y_{11}^2 y_{22}^2 + y_{11}^2 y_{32}^2) \sin[2(\phi_{32} - \phi_{31})] \right. \\ &\quad \left. + y_{22}^2 y_{33}^2 \frac{M_1}{M_3} \sin(2\phi_{32}) - y_{11}^2 y_{33}^2 \frac{M_2}{M_3} \sin(2\phi_{31}) \right]. \end{aligned} \quad (45)$$

From Table I and Eqs. (45) it is clear that we can have both $\varepsilon_1 \neq 0$ and $\mathcal{J}_{CP} \neq 0$ with a single non-vanishing phase.

On the relative sign between Y_B and $\mathcal{J}_{CP}$

It has recently been pointed out in [24] that the relative sign between CP violation and the baryon asymmetry can be predicted in a specific class of models. In our framework, it is clear from Table I and Eqs. (45) that the relative sign between the low-energy invariant $\mathcal{J}_{CP}$

TABLE II: Relative sign between the CP asymmetries $\varepsilon_{1,2}$ and the low-energy CP invariant $\mathcal{J}_{CP}$ for textures I and II considering the different heavy Majorana mass regimes. We assume $M_1, M_2 \ll M_3$ and $\phi_{32} = 0$. The parameter $x \simeq 0.4$ corresponds to the zero of the loop function $f(x)$ defined in Eq. (15).

	$\frac{M_1}{M_2} < x$	$x < \frac{M_1}{M_2} < 1$	$\frac{M_2}{M_1} < x$	$x < \frac{M_2}{M_1} < 1$
$\text{sgn}(\varepsilon_1^{\text{I}} \cdot \mathcal{J}_{CP}^{\text{I}})$	+	+	+	−
$\text{sgn}(\varepsilon_2^{\text{I}} \cdot \mathcal{J}_{CP}^{\text{I}})$	−	+	−	−
$\text{sgn}(\varepsilon_1^{\text{II}} \cdot \mathcal{J}_{CP}^{\text{II}})$	−	−	−	+
$\text{sgn}(\varepsilon_2^{\text{II}} \cdot \mathcal{J}_{CP}^{\text{II}})$	+	−	+	+

and the CP asymmetries ε_j cannot be predicted without specifying the values of the heavy neutrino masses M_i and the Dirac Yukawa couplings⁹. This is mainly due to the fact that at least one of these quantities depends on both phases appearing in the Dirac neutrino Yukawa coupling matrix for textures I and II. Therefore, in order to establish a direct connection between the sign of the CP asymmetries ε_j and the low-energy CP invariant $\mathcal{J}_{CP}$ further assumptions are required. For instance, considering the case $\phi_{32} = 0$ and $M_1, M_2 \ll M_3$ so that the terms proportional to f_{31} can be safely neglected in $\varepsilon_1^{\text{II}}$ (see Table I), we obtain from Eqs. (45) and Table I the relative signs given in Table II.

It is well known that in the case of hierarchical heavy neutrinos, the sign of Y_B is fixed by the sign of the CP asymmetry generated in the decay of the lightest heavy neutrino, let us say ε_L . For Y_B to be positive, as required by the observations, one must have $\varepsilon_L < 0$. Thus, the measurement of the sign of CP violation at low-energy could in principle exclude some of the heavy Majorana neutrino mass regimes presented in Table II. As an example, let us assume that $\mathcal{J}_{CP} < 0$. In this case, the requirement $\varepsilon_L < 0$ would exclude the mass regime $M_2/M_1 < x$ for texture I and $M_1/M_2 < x$ for texture II.

Finally, it is interesting to note that the Dirac neutrino Yukawa matrices considered in [24] correspond to our textures I and II with $y_{33} = 0$ and have the remarkable feature that the number of real parameters equals the number of masses and mixing angles. In this limit the heavy Majorana neutrino N_3 completely decouples, rendering this situation

⁹ For the sake of more generality, we do not assume the mass-ordering $M_1 < M_2$ in this discussion.

phenomenologically equivalent to the two heavy neutrino case considered in [24]. Namely, it can be easily seen that for $y_{33} = 0$ the phase ϕ_{32} can be rephased away and consequently the CP asymmetries $\varepsilon_{1,2}$ and the low-energy CP invariant $\mathcal{J}_{CP}$ will only depend on the phases ϕ_{21} and ϕ_{31} for textures I and II, respectively. This means that the examples considered in [24] are special cases of our framework which is motivated by the condition that m_D does not contain any phase that would only contribute to low-energy CP violation. It is clear from Table II that, in general, even by keeping only one phase the sign of $\varepsilon_j \cdot \mathcal{J}_{CP}$ depends on the particular Yukawa texture and hierarchy between the heavy Majorana neutrinos.

VI. EXAMPLES

In this section we present some examples of the minimal textures discussed in Section V and proceed to the study of their implications to low-energy physics as well as to the computation of the baryon asymmetry Y_B through the numerical solution of the Boltzmann equations as described in the Appendix. We will only consider cases that lead to the LMA solution of the solar neutrino problem, which means that the neutrino mixing angles and the squared mass differences lie in the typical ranges:

$$\begin{aligned} 2.5 \times 10^{-5} \text{ eV}^2 &\lesssim \Delta m_{\odot}^2 \lesssim 3.4 \times 10^{-4} \text{ eV}^2 \quad , \quad 0.24 \lesssim \tan^2 \theta_{\odot} \lesssim 0.89 \, , \\ 1.4 \times 10^{-3} \text{ eV}^2 &\lesssim \Delta m_{\oplus}^2 \lesssim 6.0 \times 10^{-3} \text{ eV}^2 \quad , \quad 0.40 \lesssim \tan^2 \theta_{\oplus} \lesssim 3.0 \, , \end{aligned} \quad (46)$$

with the solar and atmospheric mixing angles being identified as $\theta_{\odot} \equiv \theta_{12}$, $\theta_{\oplus} \equiv \theta_{23}$, respectively, in the standard parametrization [13]. The θ_{13} mixing angle is at present constrained by reactor neutrino experiments to satisfy $|\sin \theta_{13}| \equiv |U_{e3}| \lesssim 0.2$ [37].

A. Hierarchical heavy Majorana neutrinos ($M_1 < M_2 \ll M_3$)

The first example is a realization of the texture I given in Table I with $\phi_{32} = 0$. The entries of the Dirac neutrino Yukawa coupling and the right-handed neutrino mass matrices are chosen to be of order:

$$Y_{\nu} = \frac{\Lambda_D}{v} \begin{pmatrix} \epsilon^6 & 0 & 0 \\ \epsilon^6 e^{i\phi_{21}} & \epsilon^4 & 0 \\ 0 & \epsilon^4 & 1 \end{pmatrix} \, , \quad D_R = \Lambda_R \text{diag}(\epsilon^{12}, \epsilon^{10}, 1) \, , \quad (47)$$

where $\epsilon < 1$ is a small parameter. For our numerical estimates we consider the typical Dirac neutrino mass scale to be $\Lambda_D \simeq 100$ GeV, which corresponds approximately to the top quark mass at the GUT scale [38]. The neutrino mass matrix $\mathcal{M}_\nu$ is then given by:

$$\mathcal{M}_\nu = \frac{\Lambda_D^2}{\epsilon^2 \Lambda_R} \begin{pmatrix} \epsilon^2 & \epsilon^2 e^{i\phi_{21}} & 0 \\ \epsilon^2 e^{i\phi_{21}} & 1 + \epsilon^2 e^{2i\phi_{21}} & 1 \\ 0 & 1 & 1 + \epsilon^2 \end{pmatrix}. \quad (48)$$

In this particular case, considering for the moment $\phi_{21} = 0$, one gets¹⁰:

$$m_1 \simeq \frac{\Lambda_D^2}{2\Lambda_R} (2 - \sqrt{2}) \quad , \quad m_2 \simeq \frac{\Lambda_D^2}{2\Lambda_R} (2 + \sqrt{2}) \quad , \quad m_3 \simeq \frac{\Lambda_D^2}{\epsilon^2 \Lambda_R} (2 + \epsilon^2), \quad (49)$$

leading to

$$\Delta m_{21}^2 \simeq \frac{2\sqrt{2}\Lambda_D^4}{\Lambda_R^2} \quad , \quad \Delta m_{32}^2 \simeq \frac{4\Lambda_D^4}{\epsilon^4 \Lambda_R^2}. \quad (50)$$

The requirement

$$\left. \frac{\Delta m_{\odot}^2}{\Delta m_{\oplus}^2} \right|_{\text{LMA}} = \frac{\Delta m_{21}^2}{\Delta m_{32}^2} \simeq \frac{\epsilon^4}{\sqrt{2}}, \quad (51)$$

forces ϵ to be in the range $0.3 \lesssim \epsilon \lesssim 0.7$. For the neutrino mixing angles we have:

$$\tan^2 \theta_{12} \simeq 1 - \frac{\epsilon^2}{2\sqrt{2}} \quad , \quad \tan^2 \theta_{23} \simeq 1 \quad , \quad |U_{e3}| \simeq \frac{\epsilon^2}{2\sqrt{2}}, \quad (52)$$

which are in good agreement with the data taking into account the range of ϵ . The matrix V_L which corresponds to the left-handed rotation involved in the diagonalization of m_D can be viewed as the equivalent in the leptonic sector of the quark mixing matrix V_{CKM} . This matrix is obtained by diagonalizing the Hermitian matrix $Y_\nu Y_\nu^\dagger$,

$$V_L \simeq \begin{pmatrix} 1 - \epsilon^8/2 & \epsilon^4 - \epsilon^{12}/2 & 0 \\ -\epsilon^4 + \epsilon^{12}/2 & 1 - \epsilon^8/2 & \epsilon^8 \\ \epsilon^{12} & -\epsilon^8 & 1 \end{pmatrix}. \quad (53)$$

This is in fact an interesting result in the sense that we are in the presence of a typical *large mixing out of small mixing* situation, where the large neutrino mixing is generated through

¹⁰ We have checked that the light neutrino masses m_i and mixing angles θ_{ij} are not sensitive to the phase ϕ_{21} .

the seesaw mechanism [29, 39]. The scale Λ_R , or equivalently the mass M_3 of the heaviest Majorana neutrino, is determined by requiring Δm_{32}^2 to be in the experimental range given in (46). We find

$$M_3 = \Lambda_R \simeq \frac{2 \Lambda_D^2}{\epsilon^2 \sqrt{\Delta m_{\oplus}^2}} \simeq \frac{2^{5/4} \Lambda_D^2}{\sqrt{\Delta m_{\odot}^2}} \simeq 8 \times 10^{15} \text{ GeV}, \quad (54)$$

and Eq. (47) implies $M_1 \simeq 1.3 \times 10^{11} \text{ GeV}$ and $M_2 \simeq 8.4 \times 10^{12} \text{ GeV}$, for $\epsilon = 0.4$.

From the expression of the CP asymmetry ε_1^I given in Table I, we obtain

$$\varepsilon_1 = \frac{3 y_{21}^2 y_{22}^2 \sin(2 \phi_{21})}{16 \pi (y_{11}^2 + y_{21}^2)} \frac{M_1}{M_2} \simeq \frac{3 \Lambda_D^2 \epsilon^{10}}{32 \pi v^2} \sin(2 \phi_{21}) \simeq 10^{-6} \sin(2 \phi_{21}), \quad (55)$$

which is maximized for $\phi_{21} = \pi/4$. The value of the CP invariant $\mathcal{J}_{CP}$ can be computed from the expression

$$\mathcal{J}_{CP} = \frac{y_{11}^2 y_{21}^2 y_{32}^2 y_{31}^2 v^{12}}{M_1^3 M_2^3 \Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2} \left[y_{21}^2 y_{32}^2 + y_{11}^2 (y_{22}^2 + y_{32}^2) + y_{33}^2 (y_{11}^2 + y_{21}^2) \frac{M_2}{M_3} \right] \sin(2 \phi_{21}), \quad (56)$$

which is obtained from Eq. (45) setting $\phi_{32} = 0$. The result is straightforward in this case

$$\mathcal{J}_{CP} \simeq \frac{3 \epsilon^2 \sin(2 \phi_{21})}{32 \sqrt{2}} \simeq 1.1 \times 10^{-2} \sin(2 \phi_{21}), \quad (57)$$

rendering visible CP-violating effects in the next generation of neutrino factories for $\phi_{21} \simeq \pi/4$. It is interesting to note from Eqs. (55) and (57) that the dependence of ε_1 and $\mathcal{J}_{CP}$ on the phase ϕ_{21} is such that both quantities are simultaneously maximized. Notice also that ε_1 and $\mathcal{J}_{CP}$ have the same sign, as it should be according to Table II.

All the results presented above are confirmed by the full numerical computation presented in Fig. 1, where we have randomly included $\mathcal{O}(1)$ coefficients¹¹ for the non-vanishing entries of Y_ν and taken $\phi_{21} = \pi/4$. The spreading of the points in the figure, due to the random variation of the coefficients, shows that the textures are stable under these perturbations.

In order to compute the value of the baryon asymmetry we proceed to the numerical solution of the Boltzmann equations as described in the Appendix, taking the initial conditions: $Y_{N_i}^0 = 0$, $Y_{B-L}^0 = 0$. The results are presented in Fig. 2 where we have plotted Y_{N_1} , Y_{N_2} and Y_B as functions of the parameter $z = M_1/T$ for given values of the CP asymmetries, heavy

¹¹ For illustration, we have taken the $\mathcal{O}(1)$ coefficients in the range $[0.9, 1.3]$. We notice however that the results are not too sensitive to slight variations of this range.




FIG. 1: Results of the numerical computation for the textures presented in Eq. (47) for Y_ν and M_R with the CP-violating phase $\phi_{21} = \pi/4$. The heavy Majorana neutrino masses M_i and the CP asymmetries $\varepsilon_1, \varepsilon_2$ are plotted as functions of the texture parameter ϵ . The dependence of $|U_{e3}|$ and $\mathcal{J}_{CP}$ on ϵ is also shown. The dotted areas satisfy the neutrino constraints (46) for the LMA solar solution.

Majorana neutrino masses and the lightest neutrino mass. The predicted value for the final baryon asymmetry is $Y_B \simeq 9 \times 10^{-11}$, which is inside the observational range (1).

Our next example is a particular case of the texture II presented in Table I with $\phi_{32} = 0$. The Dirac neutrino Yukawa coupling and heavy neutrino mass matrices are chosen in this case to be of order

$$Y_\nu = \frac{\Lambda_D}{v} \begin{pmatrix} \epsilon^5 & 0 & 0 \\ 0 & \epsilon^3 & 0 \\ \epsilon^5 e^{i\phi_{31}} & \epsilon^3 & 1 \end{pmatrix}, \quad D_R = \Lambda_R \text{diag}(\epsilon^{10}, \epsilon^8, 1), \quad (58)$$




FIG. 2: Results for the distributions Y_{N_1} , Y_{N_2} and for the baryon asymmetry Y_B as functions of $z = M_1/T$, obtained from the numerical solution of the Boltzmann equations. The plot refers to the texture for Y_ν considered in Eq. (47) with $\phi_{21} = \pi/4$ and $\epsilon = 0.5$. The values of the heavy Majorana neutrino masses M_i , the CP asymmetries $\epsilon_{1,2}$ and the lightest neutrino mass m_1 are consistent with the ones presented in Fig. 1 for this value of ϵ . The value predicted for the final baryon asymmetry is $Y_B \simeq 9 \times 10^{-11}$.

leading to the following light neutrino mass matrix

$$\mathcal{M}_\nu = \frac{\Lambda_D^2}{\epsilon^2 \Lambda_R} \begin{pmatrix} \epsilon^2 & 0 & \epsilon^2 e^{i\phi_{31}} \\ 0 & 1 & 1 \\ \epsilon^2 e^{i\phi_{31}} & 1 & 1 + \epsilon^2(1 + e^{2i\phi_{31}}) \end{pmatrix}. \quad (59)$$

The predictions for the Δm_{ij}^2 's and neutrino mixing angles are similar to the ones given in Eqs. (50) and (52). Moreover, from Eqs. (54) and (58) and for $\epsilon = 0.4$, we get the following heavy Majorana neutrino masses: $M_1 \simeq 9 \times 10^{11} \text{ GeV}$, $M_2 \simeq 6 \times 10^{12} \text{ GeV}$, $M_3 \simeq 8 \times 10^{15} \text{ GeV}$. The left-handed matrix V_L is given in this case by:

$$V_L \simeq \begin{pmatrix} -1 & -\epsilon^{10} & \epsilon^{10} \\ -\epsilon^{10} & 1 & \epsilon^6 \\ \epsilon^{10} & -\epsilon^6 & 1 \end{pmatrix}. \quad (60)$$




FIG. 3: The same as in Fig. 1 for the textures considered in Eq. (58) and $\phi_{31} = \pi/4$.

Finally, the CP asymmetry can be obtained from the expression of $\varepsilon_1^{\text{II}}$ presented in Table I with $\phi_{32} = 0$. Taking into account the form of Y_ν and D_R as in Eqs. (58), we obtain

$$\varepsilon_1 = \frac{3 y_{31}^2 \sin(2\phi_{31})}{16 \pi (y_{11}^2 + y_{31}^2)} \left(y_{32}^2 \frac{M_1}{M_2} + y_{33}^2 \frac{M_1}{M_3} \right) \simeq \frac{3 \Lambda_D^2 \epsilon^8}{32 \pi v^2} \sin(2\phi_{31}) \simeq 6.5 \times 10^{-6} \sin(2\phi_{31}), \quad (61)$$

where the last estimate has been obtained assuming $\epsilon = 0.4$. It can also be shown that the CP invariant $\mathcal{J}_{CP}$ reads (see Eq. (45)),

$$\mathcal{J}_{CP} = -\frac{y_{11}^2 y_{22}^2 y_{31}^2 y_{32}^2 v^{12}}{M_1^3 M_2^3 \Delta m_{21}^2 \Delta m_{31}^2 \Delta m_{32}^2} \left[y_{22}^2 y_{31}^2 + y_{11}^2 (y_{22}^2 + y_{32}^2) + y_{11}^2 y_{33}^2 \frac{M_2}{M_3} \right] \sin(2\phi_{31}), \quad (62)$$

which leads to the approximate result

$$\mathcal{J}_{CP} \simeq -\frac{3 \epsilon^2 \sin(2\phi_{31})}{32 \sqrt{2}} \simeq -1.1 \times 10^{-2} \sin(2\phi_{31}). \quad (63)$$

Notice that ε_1 and $\mathcal{J}_{CP}$ have opposite signs in this case, once again in agreement with Table II.

In Fig. 3 we present the same numerical analysis as in Fig. 1, but for the case where Y_ν and D_R are defined through Eqs. (58). We find good agreement between our approximate




FIG. 4: The same as in Fig. 2 for the example considered in Eq. (58) with $\phi_{31} = \pi/4$ and $\epsilon = 0.4$. The values of the heavy Majorana neutrino masses M_i , the CP asymmetries $\epsilon_{1,2}$ and the lightest neutrino mass m_1 are consistent with the ones presented in Fig. 3 for this value of ϵ . The predicted value for the final baryon asymmetry is $Y_B \simeq 8 \times 10^{-11}$.

analytical results and the numerical ones. The integration of the Boltzmann equations is plotted in Fig. 4. The value for the final baryon asymmetry is $Y_B \simeq 8 \times 10^{-11}$.

B. Two-fold quasi-degeneracy ($M_1 \simeq M_2 \ll M_3$)

As an example, let us consider the texture of type I given in Table I and assume that $M_1 \simeq M_2 \ll M_3$. The Hermitian matrix $H_\Delta = Y_\Delta^\dagger Y_\Delta$ is given by

$$H_\Delta = \begin{pmatrix} y_{11}^2 + y_{21}^2 & y_{21} y_{22} e^{-i\phi_{21}} & 0 \\ y_{21} y_{22} e^{i\phi_{21}} & y_{22}^2 + y_{32}^2 & y_{32} y_{33} e^{-i\phi_{32}} \\ 0 & y_{32} y_{33} e^{i\phi_{32}} & y_{33}^2 \end{pmatrix}. \quad (64)$$

Taking into account the requirement of Eq. (20) we can get the range of validity of the parameter δ_N :

$$\delta_N \gg \frac{1}{8\pi} \max \{ y_{11}^2 + y_{21}^2, y_{22}^2 + y_{32}^2 \}. \quad (65)$$

Using Eqs. (23), the CP asymmetries are given in this case by

$$\varepsilon_1 \simeq \frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21})}{16\pi \delta_N (y_{11}^2 + y_{21}^2)} , \quad \varepsilon_2 \simeq \frac{1}{16\pi} \left[\frac{y_{21}^2 y_{22}^2 \sin(2\phi_{21})}{\delta_N (y_{22}^2 + y_{32}^2)} + \frac{3 y_{32}^2 y_{33}^2}{y_{22}^2 + y_{32}^2} \frac{M_2}{M_3} \sin(2\phi_{32}) \right] . \quad (66)$$

Let us now consider the following simple realization of Y_ν and D_R :

$$Y_\nu = \frac{\Lambda_D}{v} \begin{pmatrix} \epsilon^{10} & 0 & 0 \\ \epsilon^{10} e^{i\phi_{21}} & \epsilon^9 & 0 \\ 0 & \epsilon^9 & 1 \end{pmatrix} , \quad D_R = \Lambda_R \text{diag}(\epsilon^{20}, (1 + \delta_N) \epsilon^{20}, 1) , \quad (67)$$

with $\delta_N \ll 1$. From Eq. (65) we get in this case for $\epsilon = 0.3$:

$$\delta_N \gg \frac{\Lambda_D^2 \epsilon^{18}}{4\pi v^2} \simeq 10^{-11} . \quad (68)$$

The effective neutrino mass matrix is now given by

$$\mathcal{M}_\nu = \frac{\Lambda_D^2}{\epsilon^2 (1 + \delta_N) \Lambda_R} \begin{pmatrix} \epsilon^2 (1 + \delta_N) & \epsilon^2 (1 + \delta_N) e^{i\phi_{21}} & 0 \\ \epsilon^2 (1 + \delta_N) e^{i\phi_{21}} & 1 + \epsilon^2 (1 + \delta_N) e^{2i\phi_{21}} & 1 \\ 0 & 1 & 1 + \epsilon^2 (1 + \delta_N) \end{pmatrix} . \quad (69)$$

In the limit $\delta_N \rightarrow 0$, Eq. (48) is recovered. Since $\delta_N \ll 1$ the results for the neutrino masses and mixing parameters are practically the same as the ones given in Eqs. (49)-(52). The same is expected for the estimate of the CP invariant $\mathcal{J}_{CP}$ defined in Eq. (66). The main differences reside obviously in the heavy Majorana neutrino mass spectrum and on the values of the CP asymmetries, since now the relation $M_1 \ll M_2$ is no longer valid. From Eq. (23) we obtain

$$\begin{aligned} \varepsilon_1 &\simeq \frac{\epsilon^{18} \Lambda_D^2 \sin(2\phi_{21})}{32\pi \delta_N v^2} \simeq \frac{1.3 \times 10^{-12}}{\delta_N} \sin(2\phi_{21}) , \\ \varepsilon_2 &\simeq \frac{\epsilon^{20} \Lambda_D^2 \sin(2\phi_{21})}{32\pi \delta_N v^2} \simeq \frac{1.1 \times 10^{-13}}{\delta_N} \sin(2\phi_{21}) . \end{aligned} \quad (70)$$

The results of the numerical integration of the Boltzmann equations for this case are presented in Fig. 5. A realistic value for Y_B is also generated in this case.

Before we end this section, it is worthwhile to comment on the possible effects of quantum corrections to the effective neutrino mass matrix $\mathcal{M}_\nu$. This discussion turns out to be relevant since in the examples considered above we have taken the effective neutrino mass matrix at the heavy neutrino decoupling scale. Although a detailed treatment would require




FIG. 5: The same as Figs. 2 and 4 for the quasi-degenerate case $M_1 \simeq M_2$ considered in Eq. (67) with $\phi_{21} = \pi/4$, $\epsilon = 0.3$ and $\delta_N = 5 \times 10^{-7}$. The predicted value of the baryon asymmetry is $Y_B \simeq 10^{-10}$.

a renormalization group analysis for the effective neutrino mass operator, one can employ the simple analytical treatment considered by many authors in the literature (see for example Refs. [22, 40]). Following this, we recall that the effective neutrino mass matrix at the electroweak scale m_Z can be related to the one at the heavy neutrino decoupling scale Λ_R as

$$\mathcal{M}_\nu(m_Z) \propto \text{diag}(1, 1, 1 + \epsilon_\tau) \mathcal{M}_\nu(\Lambda_R) \text{diag}(1, 1, 1 + \epsilon_\tau), \quad (71)$$

where, neglecting the running of the charged lepton Yukawa couplings,

$$\epsilon_\tau \simeq \frac{3 y_\tau^2}{32 \pi^2} \ln \left(\frac{\Lambda_R}{m_Z} \right). \quad (72)$$

Taking $y_\tau = m_\tau/v$ and $\Lambda_R \lesssim 10^{16}$ GeV, we obtain $\epsilon_\tau \lesssim 10^{-5}$. Considering $\mathcal{M}_\nu(\Lambda_R)$ as given in Eq. (48) we would get from Eq. (71):

$$\begin{aligned} \frac{\Delta m_{21}^2}{\Delta m_{32}^2} &\simeq \frac{(1 - \epsilon_\tau) \epsilon^4}{\sqrt{2}} \quad , \quad \tan^2 \theta_{12} \simeq 1 - \frac{\epsilon^2}{2\sqrt{2}} - \sqrt{2} \epsilon_\tau \quad , \\ \tan^2 \theta_{23} &\simeq 1 - 2 \epsilon_\tau \quad , \quad |U_{e3}| \simeq \frac{\epsilon^2}{2\sqrt{2}} \left(1 - \frac{3}{2} \epsilon_\tau \right). \end{aligned} \quad (73)$$

Since $\epsilon_\tau \ll 1$, the results given by Eqs. (51) and (52) are not significantly altered. The same conclusions are drawn for the example considered in Eq. (59). Thus, we conclude that the effects of quantum corrections due to the renormalization of the neutrino mass operator can be safely neglected in our case.

VII. CONCLUDING REMARKS

We have analyzed, in the context of the minimal seesaw mechanism, the link between leptogenesis and CP violation at low energies. In particular, it was shown that, in order to present a thorough discussion on this question, it is convenient to work in the WB where both the charged lepton and right-handed Majorana neutrino mass matrices are diagonal and real, and to write, without loss of generality, the Dirac neutrino mass matrix as the product of a unitary matrix and a lower triangular matrix. From the analysis of the phases that contribute to leptogenesis and low-energy CP violation, we have identified a necessary condition which is required in order to establish a link between these two phenomena. We have studied a class of models which satisfy the above necessary condition in the simplest way, namely those where the Dirac neutrino mass matrix is of the triangular form. By choosing this structure the number of physical parameter in the theory is reduced then enhancing its predictability. In this case there are only three CP-violating phases which contribute both to leptogenesis and CP violation at low energies. We have then studied the minimal scenarios where a correct value of the baryon-to-entropy ratio can be generated, while accounting for all the low-energy neutrino data in the context of the LMA solution. Moreover, the examples considered in Section VI predict the existence of low-energy CP-violating effects within the range of sensitivity of the future long-baseline neutrino oscillation experiments. In fact, it is a remarkable feature of these scenarios that the solutions viable for leptogenesis are precisely those which maximize $\mathcal{J}_{CP}$. The question of relating the observed sign of the baryon asymmetry to the sign of the leptonic CP violation, measurable at low energies through neutrino oscillations, was also considered. Namely, we have concluded that, within the minimal scenarios presented, this relation crucially depends on the heavy Majorana neutrino mass spectrum. We remark that a full discussion of this aspect requires the computation of the BAU since, besides the prediction of the relative sign between the BAU and $\mathcal{J}_{CP}$, the determination of Y_B is of extreme importance to infer about the viability

of a given model.

Acknowledgments

We are grateful to Sacha Davidson for useful discussions. M.N.R. is grateful to P. Frampton and T. Yanagida for interesting conversations. This work was partially supported by *Fundação para a Ciência e a Tecnologia* (FCT, Portugal) through the projects CERN/FIS/43793/2001, POCTI/36288/FIS/2000, CERN/FIS/40139/2000 and CFIF - Plurianual (2/91) as well as by the RTN European Program HPRN-CT-2000-00148 (“AEFNET”). The work of R.G.F. and F.R.J. was supported by FCT under the grants SFRH/BPD/1549/2000 and PRAXISXXI/BD/18219/98, respectively.

APPENDIX: BOLTZMANN EQUATIONS

The computation of the cosmological baryon asymmetry Y_B involves the solution of the full set of Boltzmann equations which governs the time evolution of the right-handed neutrino number densities Y_{N_j} and the generated lepton asymmetry Y_L . These quantities depend not only on the physics occurring in the thermal bath but also on the universe expansion. In the SM framework extended with heavy right-handed neutrinos the physical processes relevant to the generation of the baryon asymmetry are typically the N_i decays and inverse decays into Higgs bosons and leptons, the $\Delta L = 1$ scatterings involving the top quark and $\Delta L = 2$ scatterings with virtual N_i and $N_i - N_j$ scatterings [41, 42]. The initially produced lepton asymmetry Y_L is converted into a net baryon asymmetry Y_B through the $(B + L)$ -violating sphaleron processes. One finds the relation

$$Y_B = \xi Y_{B-L} = \frac{\xi}{\xi - 1} Y_L. \quad (\text{A.1})$$

For the SM with three heavy Majorana neutrinos $\xi \simeq 1/3$ and therefore $Y_B \simeq Y_{B-L}/3 \simeq -Y_L/2$.

The Boltzmann equations for the N_i number densities and the Y_{B-L} asymmetry in terms of the dimensionless parameter $z = M_1/T$, where T is the temperature, can be written in

the form:

$$\begin{aligned}\frac{dY_{N_j}}{dz} &= -\frac{z}{Hs(z)} \left[\left(\gamma_j^D + \gamma_j^Q \right) \left(\frac{Y_{N_j}}{Y_{N_j}^{\text{eq}}} - 1 \right) + \sum_{i=1}^3 \gamma_{ij}^{NN} \left(\frac{Y_{N_i} Y_{N_j}}{Y_{N_i}^{\text{eq}} Y_{N_j}^{\text{eq}}} - 1 \right) \right] , \\ \frac{dY_{B-L}}{dz} &= -\frac{z}{Hs(z)} \left[\sum_{j=1}^3 \varepsilon_j \gamma_j^D \left(\frac{Y_{N_j}}{Y_{N_j}^{\text{eq}}} - 1 \right) + \gamma^W \frac{Y_{B-L}}{Y_{\ell}^{\text{eq}}} \right] ,\end{aligned}\quad (\text{A.2})$$

where H is the Hubble parameter evaluated at $z = 1$ and $s(z)$ is the entropy density given by

$$H = \sqrt{\frac{4\pi^3 g_*}{45}} \frac{M_1^2}{M_P} , \quad s(z) = \frac{2\pi^2 g_*}{45} \frac{M_1^3}{z^3} , \quad (\text{A.3})$$

respectively. Here, $g_* \simeq 106.75$ is the effective number of relativistic degrees of freedom and $M_P \simeq 1.2 \times 10^{19}$ GeV is the Planck mass. The equilibrium number density of a particle i with mass m_i is given by

$$Y_i^{\text{eq}}(z) = \frac{45}{4\pi^4} \frac{g_i}{g_*} \left(\frac{m_i}{M_1} \right)^2 z^2 K_2 \left(\frac{m_i z}{M_1} \right) , \quad (\text{A.4})$$

where g_i denote the internal degrees of freedom of the corresponding particle ($g_{N_i} = 2$, $g_{\ell} = 4$) and $K_n(x)$ are the modified Bessel functions. The γ 's are the reaction densities for the different processes. For the decays one has

$$\gamma_j^D = \frac{M_1 M_j^3}{8\pi^3 z} (Y_{\nu}^{\dagger} Y_{\nu})_{jj} K_1 \left(\frac{z M_j}{M_1} \right) . \quad (\text{A.5})$$

The reaction densities for the $\Delta L = 1$ processes with the top quark and for the $\Delta L = 2$ scatterings $N_i - N_j$ can be written in the following way:

$$\gamma_j^Q = 2\gamma_{t_j}^{(1)} + 4\gamma_{t_j}^{(2)} , \quad \gamma_{ij}^{NN} = \gamma_{N_i N_j}^{(1)} + \gamma_{N_i N_j}^{(2)} , \quad (\text{A.6})$$

respectively. Finally, γ^W accounts for the washout processes:

$$\gamma^W = \sum_{j=1}^3 \left(\frac{1}{2} \gamma_j^D + \frac{Y_{N_j}}{Y_{N_j}^{\text{eq}}} \gamma_{t_j}^{(1)} + 2\gamma_{t_j}^{(2)} \right) + 2\gamma_N^{(1)} + 2\gamma_N^{(2)} . \quad (\text{A.7})$$

Each of the above reaction densities can be computed through the corresponding reduced cross section $\hat{\sigma}^{(i)}(x)$:

$$\gamma^{(i)}(z) = \frac{M_1^4}{64\pi^4 z} \int_{(m_a^2 + m_b^2)/M_1^2}^{\infty} \hat{\sigma}^{(i)}(x) \sqrt{x} K_1(z\sqrt{x}) dx , \quad (\text{A.8})$$

where $x = s/M_1^2$, with s being the center-of-mass energy squared and $m_{a,b}$ are the masses of the initial particle states. All the relevant reduced cross sections are summarized below. For a more detailed presentation the reader is addressed to Ref. [42].

We write the reduced cross sections as functions of the parameter x , the Hermitian matrix $H_\nu = Y_\nu^\dagger Y_\nu$ and the quantities a_j and c_j defined as

$$a_j = \left(\frac{M_j}{M_1}\right)^2, \quad c_j = \left(\frac{\Gamma_j}{M_1}\right)^2 = \frac{a_j (H_\nu)_{jj}^2}{64 \pi^2}, \quad (\text{A.9})$$

where Γ_j is the decay rate defined in Eq. (19).

For the $\Delta L = 1$ processes involving interactions with quarks we have

$$\hat{\sigma}_{t_j}^{(1)}(x) \equiv \hat{\sigma}(N_j + \ell \leftrightarrow q + \bar{u}) = 3 \alpha_u (H_\nu)_{jj} \left(\frac{x - a_j}{x}\right)^2, \quad (\text{A.10})$$

$$\begin{aligned} \hat{\sigma}_{t_j}^{(2)}(x) &\equiv \hat{\sigma}(N_j + u \leftrightarrow \bar{\ell} + q) = \hat{\sigma}(N_j + \bar{q} \leftrightarrow \bar{\ell} + \bar{u}) \\ &= 3 \alpha_u (H_\nu)_{jj} \frac{x - a_j}{x} \left[1 + \frac{a_h - a_j}{x - a_j + a_h} + \frac{a_j - 2a_h}{x - a_j} \ln \left(\frac{x - a_j + a_h}{a_h} \right) \right], \end{aligned} \quad (\text{A.11})$$

where a_h (introduced to regularize the infrared divergencies) and α_u are given by

$$a_h = \left(\frac{\mu}{M_1}\right)^2, \quad \alpha_u = \frac{\text{Tr}(Y_u^\dagger Y_u)}{4\pi} \simeq \frac{m_t^2}{4\pi v^2}, \quad (\text{A.12})$$

respectively. The mass parameter μ is chosen to be $\mu = 800$ GeV as in Refs. [41, 42].

The reduced cross sections relative to the $N_i - N_j$ scatterings are:

$$\begin{aligned} \hat{\sigma}_{N_i N_j}^{(1)} &\equiv \hat{\sigma}(N_i + N_j \leftrightarrow \ell + \bar{\ell}) \\ &= \frac{1}{2\pi} \left\{ (H_\nu)_{ii} (H_\nu)_{jj} \left[\frac{\sqrt{\lambda_{ij}}}{x} + \frac{a_i + a_j}{2x} L_{ij} \right] - \text{Re} [(H_\nu)_{ji}^2] \frac{\sqrt{a_i a_j}}{x - a_i - a_j} L_{ij} \right\}, \end{aligned} \quad (\text{A.13})$$

$$\begin{aligned} \hat{\sigma}_{N_i N_j}^{(2)} &\equiv \hat{\sigma}(N_i + N_j \leftrightarrow \phi + \phi^\dagger) \\ &= \frac{1}{2\pi} \left\{ |(H_\nu)_{ji}|^2 \left[\frac{L_{ij}}{2} - \frac{\sqrt{\lambda_{ij}}}{x} \right] - \text{Re} [(H_\nu)_{ji}^2] \frac{\sqrt{a_i a_j} (a_i + a_j)}{x (x - a_i - a_j)} L_{ij} \right\}, \end{aligned} \quad (\text{A.14})$$

where the functions L_{ij} and λ_{ij} read as:

$$L_{ij} = 2 \ln \left(\frac{|x + \sqrt{\lambda_{ij}} - a_i - a_j|}{2 \sqrt{a_i a_j}} \right), \quad \lambda_{ij} = x^2 - 2x(a_i + a_j) + (a_i - a_j)^2. \quad (\text{A.15})$$

Since the heavy Majorana neutrinos are unstable particles, one has to replace the usual fermion propagators by the off-shell propagators $D_j(x)$ in the computation of the amplitudes

for the right-handed neutrino mediated processes:

$$\frac{1}{D_j} \equiv \frac{1}{D_j(x)} = \frac{x - a_j}{(x - a_j)^2 + a_j c_j}. \quad (\text{A.16})$$

For the $\Delta L = 2$ processes the reduced cross sections read then

$$\begin{aligned} \hat{\sigma}_N^{(1)} &\equiv \hat{\sigma}(\ell + \phi \leftrightarrow \bar{\ell} + \phi^\dagger) = \sum_{j=1}^3 (H_\nu)_{jj}^2 \mathcal{A}_{jj}^{(1)} + \sum_{\substack{n,j=1 \\ j < n}}^3 \text{Re} \left[(H_\nu)_{nj}^2 \right] \mathcal{B}_{nj}^{(1)}, \\ \hat{\sigma}_N^{(2)} &\equiv \hat{\sigma}(\ell + \ell \leftrightarrow \phi^\dagger + \phi^\dagger) = \sum_{j=1}^3 (H_\nu)_{jj}^2 \mathcal{A}_{jj}^{(2)} + \sum_{\substack{n,j=1 \\ j < n}}^3 \text{Re} \left[(H_\nu)_{nj}^2 \right] \mathcal{B}_{nj}^{(2)}, \end{aligned} \quad (\text{A.17})$$

where

$$\begin{aligned} \mathcal{A}_{jj}^{(1)} &= \frac{1}{2\pi} \left[1 + \frac{a_j}{D_j} + \frac{a_j x}{2 D_j^2} - \frac{a_j}{x} \left(1 + \frac{x + a_j}{D_j} \right) \ln \left(\frac{x + a_j}{a_j} \right) \right], \\ \mathcal{B}_{nj}^{(1)} &= \frac{\sqrt{a_n a_j}}{2\pi} \left[\frac{1}{D_j} + \frac{1}{D_n} + \frac{x}{D_j D_n} + \left(1 + \frac{a_j}{x} \right) \left(\frac{2}{a_n - a_j} - \frac{1}{D_n} \right) \ln \left(\frac{x + a_j}{a_j} \right) \right. \\ &\quad \left. + \left(1 + \frac{a_n}{x} \right) \left(\frac{2}{a_j - a_n} - \frac{1}{D_j} \right) \ln \left(\frac{x + a_n}{a_n} \right) \right], \\ \mathcal{A}_{jj}^{(2)} &= \frac{1}{2\pi} \left[\frac{x}{x + a_j} + \frac{a_j}{x + 2a_j} \ln \left(\frac{x + a_j}{a_j} \right) \right], \\ \mathcal{B}_{nj}^{(2)} &= \frac{\sqrt{a_n a_j}}{2\pi} \left\{ \frac{1}{x + a_n + a_j} \ln \left[\frac{(x + a_j)(x + a_n)}{a_j a_n} \right] + \frac{2}{a_n - a_j} \ln \left[\frac{a_n (x + a_j)}{a_j (x + a_n)} \right] \right\}. \end{aligned} \quad (\text{A.18})$$

In our analysis we have computed numerically the reaction densities through Eq. (A.8) and the above definitions of the reduced cross sections. Nevertheless, useful analytical approximations can be obtained for specific temperature regimes [42].

-
- [1] S. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **86**, 5651 (2001) [arXiv:hep-ex/0103032]; S. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **86**, 5656 (2001) [arXiv:hep-ex/0103033].
 - [2] Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **87**, 071301 (2001) [arXiv:nucl-ex/0106015].
 - [3] For recent reviews and a set of references on neutrino mass models see, *e.g.*, I. Masina, Int. J. Mod. Phys. A **16**, 5101 (2001) [arXiv:hep-ph/0107220]; G. Altarelli and F. Feruglio, arXiv:hep-ph/0206077; S. F. King, arXiv:hep-ph/0208266.

- [4] T. Yanagida, in *Proc. of the Workshop on the Unified Theory and Baryon Number in the Universe*, ed. by O. Sawada and A. Sugamoto (KEK report 79-18, 1979), p.95, Tsukuba, Japan; M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, ed. by P. van Nieuwenhuizen and D. Z. Freedman (North Holland, Amsterdam, 1979), p.315; R. N. Mohapatra and G. Senjanovic, *Phys. Rev. Lett.* **44**, 912 (1980).
- [5] G. Jungman, M. Kamionkowski, A. Kosowsky and D. N. Spergel, *Phys. Rev. D* **54**, 1332 (1996); M. Zaldarriaga, D. N. Spergel and U. Seljak, *Astrophys. J.* **488**, 1 (1997); P. de Bernardis *et al.*, *Astrophys. J.* **564**, 559 (2002); C. Pryke, N. W. Halverson, E. M. Leitch, J. Kovac, J. E. Carlstrom, W. L. Holzapfel and M. Dragovan, *Astrophys. J.* **568**, 46 (2002).
- [6] http://map.gsfc.nasa.gov/m_mm/ms_status.html
- [7] <http://astro.estec.esa.nl/Planck/>
- [8] M. Fukugita and T. Yanagida, *Phys. Lett. B* **174**, 45 (1986).
- [9] For recent discussions on the physics potential of future long-baseline neutrino oscillation experiments see, *e.g.*, J. Burguet-Castell, M. B. Gavela, J. J. Gomez-Cadenas, P. Hernandez and O. Mena, [arXiv:hep-ph/0207080](https://arxiv.org/abs/hep-ph/0207080); M. Lindner, [arXiv:hep-ph/0209083](https://arxiv.org/abs/hep-ph/0209083), and references therein.
- [10] T. Endoh, T. Morozumi, T. Onogi and A. Purwanto, *Phys. Rev. D* **64**, 013006 (2001) [Erratum, *ibid.* **D 64**, 059904 (2001)] [[arXiv:hep-ph/0012345](https://arxiv.org/abs/hep-ph/0012345)].
- [11] G. C. Branco, L. Lavoura and M. N. Rebelo, *Phys. Lett. B* **180**, 264 (1986).
- [12] G. C. Branco, T. Morozumi, B. M. Nobre and M. N. Rebelo, *Nucl. Phys. B* **617**, 475 (2001) [[arXiv:hep-ph/0107164](https://arxiv.org/abs/hep-ph/0107164)].
- [13] K. Hagiwara *et al.* [Particle Data Group Collaboration], *Phys. Rev. D* **66**, 010001 (2002).
- [14] G. C. Branco, M. N. Rebelo and J. I. Silva-Marcos, *Phys. Rev. Lett.* **82**, 683 (1999)[[hep-ph/9810328](https://arxiv.org/abs/hep-ph/9810328)].
- [15] L. Covi, E. Roulet and F. Vissani, *Phys. Lett. B* **384**, 169 (1996) [[arXiv:hep-ph/9605319](https://arxiv.org/abs/hep-ph/9605319)]; W. Buchmüller and M. Plümacher, *Phys. Lett. B* **431**, 354 (1998) [[arXiv:hep-ph/9710460](https://arxiv.org/abs/hep-ph/9710460)].
- [16] W. Buchmüller and S. Fredenhagen, *Phys. Lett. B* **483**, 217 (2000) [[arXiv:hep-ph/0004145](https://arxiv.org/abs/hep-ph/0004145)].
- [17] S. Davidson and A. Ibarra, *Phys. Lett. B* **535**, 25 (2002) [[arXiv:hep-ph/0202239](https://arxiv.org/abs/hep-ph/0202239)].
- [18] J. R. Ellis, M. Raidal and T. Yanagida, *Phys. Lett. B* **546**, 228 (2002) [[arXiv:hep-ph/0206300](https://arxiv.org/abs/hep-ph/0206300)].
- [19] J. R. Ellis, J. E. Kim and D. V. Nanopoulos, *Phys. Lett. B* **145**, 181 (1984); J. R. Ellis, D. V. Nanopoulos and S. Sarkar, *Nucl. Phys. B* **259**, 175 (1985); T. Moroi, H. Murayama and M. Yamaguchi, *Phys. Lett. B* **303**, 289 (1993); M. Kawasaki and T. Moroi, *Prog. Theor.*

- Phys. **93**, 879 (1995) [arXiv:hep-ph/9403364]; M. Bolz, A. Brandenburg and W. Buchmüller, Nucl. Phys. B **606**, 518 (2001) [arXiv:hep-ph/0012052].
- [20] G. F. Giudice, M. Peloso, A. Riotto and I. Tkachev, JHEP **9908**, 014 (1999) [arXiv:hep-ph/9905242]; T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, Phys. Lett. B **464**, 12 (1999) [arXiv:hep-ph/9906366]; T. Asaka, K. Hamaguchi, M. Kawasaki and T. Yanagida, Phys. Rev. D **61**, 083512 (2000) [arXiv:hep-ph/9907559]; J. Garcia-Bellido and E. Ruiz Morales, Phys. Lett. B **536**, 193 (2002) [arXiv:hep-ph/0109230]; M. Fujii, K. Hamaguchi and T. Yanagida, Phys. Rev. D **65**, 115012 (2002) [arXiv:hep-ph/0202210]; R. Allahverdi and A. Mazumdar, Phys. Rev. D **67**, 023509 (2003) [arXiv:hep-ph/0208268]. L. Boubekur, S. Davidson, M. Peloso and L. Sorbo, arXiv:hep-ph/0209256.
- [21] M. Flanz, E. A. Paschos and U. Sarkar, Phys. Lett. B **345**, 248 (1995) [Erratum-ibid. B **382**, 447 (1996)] [arXiv:hep-ph/9411366]; A. Pilaftsis, Phys. Rev. D **56**, 5431 (1997) [arXiv:hep-ph/9707235]; W. Buchmüller and M. Plümacher, Phys. Lett. B **431**, 354 (1998) [arXiv:hep-ph/9710460]; A. Pilaftsis, Int. J. Mod. Phys. A **14**, 1811 (1999) [arXiv:hep-ph/9812256].
- [22] R. González Felipe and F. R. Joaquim, JHEP **0109**, 015 (2001) [arXiv:hep-ph/0106226].
- [23] G. C. Branco, R. González Felipe, F. R. Joaquim and M. N. Rebelo, Nucl. Phys. B **640**, 202 (2002) [arXiv:hep-ph/0202030].
- [24] P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. B **548**, 119 (2002) [arXiv:hep-ph/0208157].
- [25] J. A. Casas and A. Ibarra, Nucl. Phys. B **618**, 171 (2001) [arXiv:hep-ph/0103065].
- [26] S. Lavignac, I. Masina and C. A. Savoy, Nucl. Phys. B **633**, 139 (2002) [arXiv:hep-ph/0202086].
- [27] I. Masina, arXiv:hep-ph/0210125.
- [28] M. N. Rebelo, Phys. Rev. D **67**, 013008 (2003) [arXiv:hep-ph/0207236].
- [29] G. Altarelli, F. Feruglio and I. Masina, Phys. Lett. B **472**, 382 (2000) [arXiv:hep-ph/9907532].
- [30] A. Y. Smirnov, Phys. Rev. D **48**, 3264 (1993) [arXiv:hep-ph/9304205]; S. F. King, Nucl. Phys. B **562**, 57 (1999) [arXiv:hep-ph/9904210]; S. F. King, Nucl. Phys. B **576**, 85 (2000) [arXiv:hep-ph/9912492]; M. Hirsch and S. F. King, Phys. Rev. D **64**, 113005 (2001) [arXiv:hep-ph/0107014]; S. F. King, JHEP **0209**, 0411 (2002) [arXiv:hep-ph/0204360].
- [31] Lisbon-Saclay collaboration, in preparation.
- [32] J. R. Ellis and M. Raidal, Nucl. Phys. B **643**, 229 (2002) [arXiv:hep-ph/0206174]; S. Davidson and A. Ibarra, Nucl. Phys. B **648**, 345 (2003) [arXiv:hep-ph/0206304].

- [33] W. Buchmüller and D. Wyler, Phys. Lett. B **521**, 291 (2001) [arXiv:hep-ph/0108216]; F. Buccella, D. Falcone and F. Tramontano, Phys. Lett. B **524**, 241 (2002) [arXiv:hep-ph/0108172]; J. R. Ellis, J. Hisano, S. Lola and M. Raidal, Nucl. Phys. B **621**, 208 (2002) [arXiv:hep-ph/0109125]; W. Rodejohann and K. R. Balaji, Phys. Rev. D **65**, 093009 (2002) [arXiv:hep-ph/0201052]; H. B. Nielsen and Y. Takanishi, Nucl. Phys. B **636**, 305 (2002) [arXiv:hep-ph/0204027]; W. Buchmüller, P. Di Bari and M. Plümacher, Nucl. Phys. B **643**, 367 (2002) [arXiv:hep-ph/0205349]; W. Rodejohann, Phys. Lett. B **542**, 100 (2002) [arXiv:hep-ph/0207053]; T. Endoh, S. Kaneko, S. K. Kang, T. Morozumi and M. Tanimoto, Phys. Rev. Lett. **89**, 231601 (2002) [arXiv:hep-ph/0209020]. Z. Z. Xing, arXiv:hep-ph/0209066; J. C. Pati, arXiv:hep-ph/0209160; M. Raidal and A. Strumia, Phys. Lett. B **553**, 72 (2003) [arXiv:hep-ph/0210021]. T. Hambye, arXiv:hep-ph/0210048; S. Kaneko and M. Tanimoto, Phys. Lett. B **551**, 127 (2003) [arXiv:hep-ph/0210155].
- [34] M. Plümacher, Z. Phys. C **74**, 549 (1997) [arXiv:hep-ph/9604229].
- [35] A. Kageyama, S. Kaneko, N. Shimoyama and M. Tanimoto, Phys. Lett. B **538**, 96 (2002) [arXiv:hep-ph/0204291].
- [36] P. H. Frampton, S. L. Glashow and D. Marfatia, Phys. Lett. B **536**, 79 (2002) [arXiv:hep-ph/0201008].
- [37] M. Apollonio *et al.* [CHOOZ Collaboration], Phys. Lett. B **466**, 415 (1999) [arXiv:hep-ex/9907037].
- [38] H. Fusaoka and Y. Koide, Phys. Rev. D **57**, 3986 (1998) [arXiv:hep-ph/9712201].
- [39] E. K. Akhmedov, G. C. Branco, F. R. Joaquim and J. I. Silva-Marcos, Phys. Lett. B **498**, 237 (2001) [arXiv:hep-ph/0008010].
- [40] J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. B **556**, 3 (1999) [arXiv:hep-ph/9904395]; R. Barbieri, G. G. Ross and A. Strumia, JHEP **9910**, 020 (1999) [arXiv:hep-ph/9906470]; N. Haba, Y. Matsui and N. Okamura, Prog. Theor. Phys. **103**, 807 (2000) [arXiv:hep-ph/9911481].
- [41] M. A. Luty, Phys. Rev. D **45**, 455 (1992).
- [42] M. Plümacher, Nucl. Phys. B **530**, 207 (1998) [arXiv:hep-ph/9704231].